



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 11, 2024 – 09:36 PM EDT

PDB ID : 1Y14
Title : Crystal structure of yeast subcomplex of Rpb4 and Rpb7
Authors : Armache, K.-J.; Mitterweger, S.; Meinhart, A.; Cramer, P.
Deposited on : 2004-11-17
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	187	<div> <div>5%</div> <div> <div></div> <div>51%</div> <div>18%</div> <div>••</div> <div>29%</div> </div> </div>
1	C	187	<div> <div>6%</div> <div> <div></div> <div>50%</div> <div>22%</div> <div>•</div> <div>27%</div> </div> </div>
2	B	171	<div> <div>8%</div> <div> <div></div> <div>60%</div> <div>27%</div> <div>••</div> <div>10%</div> </div> </div>
2	D	171	<div> <div>4%</div> <div> <div></div> <div>61%</div> <div>32%</div> <div>•</div> <div>6%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4776 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II 32 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	133	Total	C	N	O	S	0	0	0
			1059	659	185	213	2			
1	C	136	Total	C	N	O	S	0	0	0
			1087	677	192	216	2			

- Molecule 2 is a protein called DNA-directed RNA polymerase II 19 kDa polypeptide.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	154	Total	C	N	O	S	0	0	0
			1220	790	198	224	8			
2	D	160	Total	C	N	O	S	0	0	0
			1256	811	203	234	8			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	43	Total	O	0	0
			43	43		
3	B	30	Total	O	0	0
			30	30		
3	C	37	Total	O	0	0
			37	37		
3	D	44	Total	O	0	0
			44	44		

- Molecule 1: DNA-directed RNA polymerase II 32 kDa polypeptide





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	103.65Å 114.81Å 80.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 46.73 – 2.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.30) 95.3 (46.73-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.43 (at 2.29Å)	Xtriage
Refinement program	unknown	Depositor
R, R_{free}	0.228 , 0.274 0.232 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	46.2	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 57.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4776	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/1066	0.61	0/1432
1	C	0.36	0/1095	0.61	0/1469
2	B	0.41	0/1243	0.68	0/1670
2	D	0.43	0/1280	0.69	0/1721
All	All	0.39	0/4684	0.65	0/6292

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1059	0	1083	30	0
1	C	1087	0	1116	36	0
2	B	1220	0	1239	55	0
2	D	1256	0	1272	44	0
3	A	43	0	0	0	0
3	B	30	0	0	0	0
3	C	37	0	0	1	0
3	D	44	0	0	0	0
All	All	4776	0	4710	155	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 155 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:138:THR:HG22	2:B:139:ILE:H	1.42	0.85
2:D:126:ASN:HD22	2:D:127:PRO:HA	1.44	0.82
2:B:56:ILE:O	2:B:57:GLN:HG2	1.82	0.79
2:D:13:LEU:HD21	2:D:17:PHE:HB2	1.68	0.76
2:B:138:THR:HG22	2:B:139:ILE:N	2.00	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/187 (69%)	121 (94%)	5 (4%)	3 (2%)	6	5
1	C	132/187 (71%)	126 (96%)	5 (4%)	1 (1%)	19	23
2	B	148/171 (86%)	135 (91%)	10 (7%)	3 (2%)	7	6
2	D	154/171 (90%)	146 (95%)	7 (4%)	1 (1%)	25	31
All	All	563/716 (79%)	528 (94%)	27 (5%)	8 (1%)	11	11

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	15	PRO
1	A	73	SER
2	B	134	GLU
1	C	136	GLY
1	A	138	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	119/169 (70%)	113 (95%)	6 (5%)	24	34
1	C	122/169 (72%)	118 (97%)	4 (3%)	38	53
2	B	139/152 (91%)	136 (98%)	3 (2%)	52	69
2	D	143/152 (94%)	138 (96%)	5 (4%)	36	50
All	All	523/642 (82%)	505 (97%)	18 (3%)	37	51

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	57	GLN
2	D	165	GLU
2	D	142	ARG
2	B	141	SER
2	D	10	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	131	GLN
2	D	158	HIS
1	C	137	ASN
1	C	143	ASN
1	C	200	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	133/187 (71%)	0.44	10 (7%) 14 19	28, 47, 86, 99	0
1	C	136/187 (72%)	0.37	12 (8%) 10 13	30, 51, 93, 100	0
2	B	154/171 (90%)	0.74	14 (9%) 9 12	29, 51, 92, 99	0
2	D	160/171 (93%)	0.40	7 (4%) 34 41	27, 47, 89, 99	0
All	All	583/716 (81%)	0.49	43 (7%) 14 19	27, 49, 92, 100	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	77	HIS	6.3
2	D	21	ARG	5.9
1	A	136	GLY	5.1
1	C	118	THR	5.0
1	A	47	LEU	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.