



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 3, 2025 – 03:53 PM JST

PDB ID : 8YG7  
Title : HSA Copper Indomethacin Complex  
Authors : Zhang, Z.L.; Yang, F.  
Deposited on : 2024-02-26  
Resolution : 2.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.2


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	581	<div> <div style="width: 83%;"></div> <div style="width: 15%;"></div> <div style="width: 2%;"></div> </div> <div>83% 15% .</div>
1	I	581	<div> <div style="width: 83%;"></div> <div style="width: 13%;"></div> <div style="width: 4%;"></div> </div> <div>83% 13% .</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IMN	A	602	-	-	X	-

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Albumin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	581	Total	C	N	O	S	0	0	0
			4443	2819	744	839	41			
1	I	581	Total	C	N	O	S	0	0	0
			4443	2819	744	839	41			

- Molecule 2 is {N}, {N}-dimethyl-7-propan-2-yl-3-thia-5-aza-1,6-diazonia-2<sup>3</sup>-cupratricyclo[6.4.0.0<sup>2,6</sup>]dodeca-1(12),4,6,8,10-pentaen-4-amine (three-letter code: A1D6U) (formula: C<sub>12</sub>H<sub>17</sub>CuN<sub>4</sub>S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cu	N	S	0	0
			18	12	1	4	1		
2	I	1	Total	C	Cu	N	S	0	0
			18	12	1	4	1		

- Molecule 3 is INDOMETHACIN (three-letter code: IMN) (formula: C<sub>19</sub>H<sub>16</sub>ClNO<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0
			25	19	1	1	4	
3	I	1	Total	C	Cl	N	O	0
			25	19	1	1	4	

- Molecule 4 is PALMITIC ACID (three-letter code: PLM) (formula: C<sub>16</sub>H<sub>32</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C	O	0
			13	11	2	
4	A	1	Total	C	O	0
			18	16	2	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			17	15	2		
4	A	1	Total	C	O	0	0
			18	16	2		
4	A	1	Total	C	O	0	0
			17	15	2		
4	A	1	Total	C	O	0	0
			18	16	2		
4	I	1	Total	C	O	0	0
			13	11	2		
4	I	1	Total	C	O	0	0
			18	16	2		
4	I	1	Total	C	O	0	0
			17	15	2		
4	I	1	Total	C	O	0	0
			18	16	2		
4	I	1	Total	C	O	0	0
			17	15	2		
4	I	1	Total	C	O	0	0
			18	16	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	129	Total	O	0	0
			129	129		
5	I	117	Total	O	0	0
			117	117		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Albumin



- Molecule 1: Albumin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.35Å 92.61Å 95.22Å 75.02° 89.28° 79.82°	Depositor
Resolution (Å)	91.93 – 2.09 91.93 – 2.09	Depositor EDS
% Data completeness (in resolution range)	97.3 (91.93-2.09) 87.5 (91.93-2.09)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.262 , 0.325 0.265 , 0.325	Depositor DCC
$R_{free}$ test set	3593 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.5	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 26.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.198 for h,h-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9420	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLM, IMN, A1D6U

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.69	0/4532	0.84	8/6148 (0.1%)
1	I	0.68	0/4532	0.83	8/6148 (0.1%)
All	All	0.68	0/9064	0.84	16/12296 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	I	0	1

There are no bond length outliers.

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	144	ARG	NE-CZ-NH2	-6.63	116.99	120.30
1	I	81	ARG	NE-CZ-NH1	6.02	123.31	120.30
1	A	218	ARG	NE-CZ-NH1	6.01	123.31	120.30
1	A	144	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	I	336	ARG	NE-CZ-NH2	-5.68	117.46	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	I	95	GLU	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4443	0	4193	55	0
1	I	4443	0	4193	64	0
2	A	18	0	0	0	0
2	I	18	0	0	0	0
3	A	25	0	15	10	0
3	I	25	0	15	7	0
4	A	101	0	163	5	0
4	I	101	0	163	4	0
5	A	129	0	0	10	0
5	I	117	0	0	17	0
All	All	9420	0	8742	126	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 126 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:411:TYR:N	5:A:701:HOH:O	1.97	0.98
1:A:199:LYS:HB3	3:A:602:IMN:CL	2.08	0.91
1:I:40:VAL:N	5:I:702:HOH:O	2.05	0.86
1:I:464:HIS:NE2	5:I:701:HOH:O	1.90	0.83
1:I:404:GLN:HE21	1:I:428:ARG:HA	1.42	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	579/581 (100%)	547 (94%)	27 (5%)	5 (1%)	14	11
1	I	579/581 (100%)	550 (95%)	23 (4%)	6 (1%)	13	9
All	All	1158/1162 (100%)	1097 (95%)	50 (4%)	11 (1%)	14	11

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	399	GLY
1	A	469	VAL
1	A	563	ASP
1	A	569	ALA
1	I	314	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	455/509 (89%)	430 (94%)	25 (6%)	18	16
1	I	455/509 (89%)	424 (93%)	31 (7%)	13	11
All	All	910/1018 (89%)	854 (94%)	56 (6%)	15	13

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	74	LEU
1	I	563	ASP
1	I	219	LEU
1	I	500	LYS
1	I	469	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	109	ASN
1	A	464	HIS
1	I	61	ASN
1	I	390	GLN
1	I	404	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	PLM	A	603	-	12,12,17	0.82	0	12,12,17	0.52	0
2	A1D6U	A	601	1	15,20,20	3.42	5 (33%)	18,30,30	4.02	5 (27%)
4	PLM	I	608	-	17,17,17	0.59	0	17,17,17	0.69	0
2	A1D6U	I	601	1	15,20,20	3.50	5 (33%)	18,30,30	3.97	5 (27%)
4	PLM	A	604	-	17,17,17	0.55	0	17,17,17	0.81	0
4	PLM	A	606	-	17,17,17	0.61	0	17,17,17	0.73	0
4	PLM	A	607	-	16,16,17	0.68	0	16,16,17	0.80	0
4	PLM	A	608	-	17,17,17	0.56	0	17,17,17	0.73	0
4	PLM	I	605	-	16,16,17	0.51	0	16,16,17	0.99	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	IMN	A	602	-	24,27,27	1.66	5 (20%)	29,39,39	3.10	11 (37%)
4	PLM	I	607	-	16,16,17	0.78	0	16,16,17	0.71	0
4	PLM	I	604	-	17,17,17	0.65	0	17,17,17	0.57	0
4	PLM	I	603	-	12,12,17	0.75	0	12,12,17	0.88	1 (8%)
3	IMN	I	602	-	24,27,27	1.76	5 (20%)	29,39,39	4.05	13 (44%)
4	PLM	I	606	-	17,17,17	0.45	0	17,17,17	0.93	1 (5%)
4	PLM	A	605	-	16,16,17	0.47	0	16,16,17	1.06	2 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PLM	A	603	-	-	7/10/10/15	-
2	A1D6U	A	601	1	-	4/8/32/32	0/3/3/3
4	PLM	I	608	-	-	7/15/15/15	-
2	A1D6U	I	601	1	-	4/8/32/32	0/3/3/3
4	PLM	A	604	-	-	10/15/15/15	-
4	PLM	A	606	-	-	7/15/15/15	-
4	PLM	A	607	-	-	6/14/14/15	-
4	PLM	A	608	-	-	6/15/15/15	-
4	PLM	I	605	-	-	6/14/14/15	-
3	IMN	A	602	-	-	4/10/14/14	0/3/3/3
4	PLM	I	607	-	-	8/14/14/15	-
4	PLM	I	604	-	-	8/15/15/15	-
4	PLM	I	603	-	-	5/10/10/15	-
3	IMN	I	602	-	-	5/10/14/14	0/3/3/3
4	PLM	I	606	-	-	9/15/15/15	-
4	PLM	A	605	-	-	9/14/14/15	-

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	601	A1D6U	C00J-S003	8.50	1.82	1.73
2	A	601	A1D6U	C00J-N00E	8.07	1.48	1.35
2	I	601	A1D6U	C00J-N00E	8.04	1.48	1.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	A1D6U	C00J-S003	7.98	1.82	1.73
3	I	602	IMN	C10-C9	4.96	1.58	1.50

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	A1D6U	S003-C00J-N00C	-15.51	110.22	124.85
2	I	601	A1D6U	S003-C00J-N00C	-15.19	110.53	124.85
3	I	602	IMN	C7-C17-C18	12.46	129.15	114.18
3	I	602	IMN	C10-C9-N	11.28	131.12	117.95
3	A	602	IMN	C16-C8-N	8.20	132.71	122.37

There are no chirality outliers.

5 of 105 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	A1D6U	S003-C00J-N00E-C00Z
2	I	601	A1D6U	S003-C00J-N00E-C00Z
2	I	601	A1D6U	N00C-C00J-N00E-C00Z
3	I	602	IMN	C2-C3-O-C6
3	I	602	IMN	C4-C3-O-C6


There are no ring outliers.



7 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	603	PLM	1	0
4	A	604	PLM	1	0
4	A	606	PLM	3	0
3	A	602	IMN	10	0
4	I	604	PLM	1	0
3	I	602	IMN	7	0
4	I	606	PLM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the


average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## Ligand A1D6U I 601



## Ligand PLM A 604









## Ligand PLM I 607




## Ligand PLM I 604




## Ligand PLM I 603




## Ligand IMN I 602



## Ligand PLM I 606





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	581/581 (100%)	-1.20	0 100 100	11, 32, 56, 73	0
1	I	581/581 (100%)	-1.14	0 100 100	12, 34, 59, 84	0
All	All	1162/1162 (100%)	-1.17	0 100 100	11, 33, 58, 84	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	A1D6U	I	601	18/18	0.97	0.09	77,91,104,114	0
3	IMN	A	602	25/25	0.97	0.09	66,73,114,129	0
3	IMN	I	602	25/25	0.97	0.07	39,74,80,91	0
4	PLM	A	603	13/18	0.97	0.06	35,39,43,44	0
4	PLM	I	604	18/18	0.97	0.06	34,39,50,53	0
4	PLM	A	604	18/18	0.98	0.06	34,40,46,46	0
4	PLM	A	605	17/18	0.98	0.06	25,45,49,50	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PLM	A	606	18/18	0.98	0.08	34,42,58,60	0
4	PLM	A	607	17/18	0.98	0.06	33,35,50,53	0
4	PLM	A	608	18/18	0.98	0.07	38,44,59,60	0
4	PLM	I	603	13/18	0.98	0.06	44,49,53,53	0
2	A1D6U	A	601	18/18	0.98	0.07	74,84,100,105	0
4	PLM	I	605	17/18	0.98	0.07	33,40,46,48	0
4	PLM	I	606	18/18	0.98	0.08	44,51,68,70	0
4	PLM	I	607	17/18	0.98	0.08	40,44,61,61	0
4	PLM	I	608	18/18	0.98	0.09	45,57,74,77	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around A1D6U I 601:**




2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around IMN A 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around IMN I 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around PLM A 603:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around PLM I 604:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around PLM A 604:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around PLM A 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around PLM A 606:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around PLM A 607:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around PLM A 608:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around PLM I 603:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around A1D6U A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around PLM I 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around PLM I 606:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around PLM I 607:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around PLM I 608:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.