



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 31, 2025 – 09:01 PM JST

PDB ID : 5YJA / pdb\_00005yja  
Title : Crystal structure of highly active BTUO mutant P287G without dehydration  
Authors : Hibi, T.; Itoh, T.; Nishiya, Y.  
Deposited on : 2017-10-09  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.42


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2328 (1.66-1.66)
Clashscore	180529	2515 (1.66-1.66)
Ramachandran outliers	177936	2475 (1.66-1.66)
Sidechain outliers	177891	2475 (1.66-1.66)
RSRZ outliers	164620	2328 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>5%</div> <div>8%</div> </div> </div>
1	B	323	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>7%</div> <div>7%</div> </div> </div>
1	C	323	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>6%</div> <div>7%</div> </div> </div>
1	D	323	<div> <div>4%</div> <div> <div></div> <div>89%</div> <div>•</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 19802 atoms, of which 9135 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Uric acid degradation bifunctional protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	298	Total	C	H	N	O	S	0	0	0
			4586	1511	2246	382	443	4			
1	B	302	Total	C	H	N	O	S	0	7	0
			4808	1568	2360	404	472	4			
1	C	299	Total	C	H	N	O	S	0	7	0
			4732	1558	2311	395	464	4			
1	D	296	Total	C	H	N	O	S	0	0	0
			4495	1491	2190	377	433	4			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	287	GLY	PRO	engineered mutation	UNP Q45697
B	287	GLY	PRO	engineered mutation	UNP Q45697
C	287	GLY	PRO	engineered mutation	UNP Q45697
D	287	GLY	PRO	engineered mutation	UNP Q45697

- Molecule 2 is 8-AZAXANTHINE (CCD ID: AZA) (formula: C<sub>4</sub>H<sub>3</sub>N<sub>5</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			14	4	3	5	2		
2	B	1	Total	C	H	N	O	0	0
			14	4	3	5	2		
2	C	1	Total	C	H	N	O	0	0
			14	4	3	5	2		
2	D	1	Total	C	H	N	O	0	0
			14	4	3	5	2		

- Molecule 3 is OXYGEN MOLECULE (CCD ID: OXY) (formula: O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O 2 2	0	0
3	B	1	Total O 2 2	0	0
3	C	1	Total O 2 2	0	0
3	D	1	Total O 2 2	0	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0

- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total K 1 1	0	0
5	C	1	Total K 1 1	0	0

- Molecule 6 is 2-METHOXYETHANOL (CCD ID: MXE) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	H	O	0	0
			13	3	8	2		
6	C	1	Total	C	H	O	0	0
			13	3	8	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	228	Total	O	0	0
			228	228		
7	B	326	Total	O	0	0
			326	326		
7	C	304	Total	O	0	0
			304	304		
7	D	221	Total	O	0	0
			221	221		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uric acid degradation bifunctional protein




- Molecule 1: Uric acid degradation bifunctional protein



- Molecule 1: Uric acid degradation bifunctional protein



- Molecule 1: Uric acid degradation bifunctional protein



PHE  
SER  
ASP  
GLU



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	133.02Å 144.25Å 70.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.68 – 1.65 39.68 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (39.68-1.65) 99.8 (39.68-1.65)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.75 (at 1.65Å)	Xtriage
Refinement program	PHENIX (1.12_2829)	Depositor
R, $R_{free}$	0.164 , 0.187 0.164 , 0.186	Depositor DCC
$R_{free}$ test set	16325 reflections (10.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.3	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 48.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	19802	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 54.11 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.8281e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OXY, AZA, OCS, SO4, MXE, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/2385	0.61	0/3236
1	B	0.49	0/2499	0.64	0/3389
1	C	0.49	0/2466	0.67	0/3346
1	D	0.43	0/2348	0.58	0/3189
All	All	0.46	0/9698	0.63	0/13160

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2340	2246	2242	9	0
1	B	2448	2360	2344	11	0
1	C	2421	2311	2309	11	0
1	D	2305	2190	2188	5	0
2	A	11	3	3	1	0
2	B	11	3	3	1	0
2	C	11	3	3	1	0
2	D	11	3	3	1	0
3	A	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
6	B	5	8	8	1	0
6	C	5	8	8	0	0
7	A	228	0	0	0	0
7	B	326	0	0	4	0
7	C	304	0	0	3	0
7	D	221	0	0	2	0
All	All	10667	9135	9111	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:234:ARG:NH2	7:C:501:HOH:O	2.27	0.68
1:B:201:ARG:HH12	2:B:401:AZA:HN3	1.42	0.68
1:C:201:ARG:HH12	2:C:401:AZA:HN3	1.41	0.66
1:D:201:ARG:HH12	2:D:401:AZA:HN3	1.44	0.65
1:A:201:ARG:HH12	2:A:401:AZA:HN3	1.42	0.65
1:B:40:ARG:NH1	7:B:501:HOH:O	2.31	0.64
1:B:9:MET:N	7:B:502:HOH:O	2.31	0.62
1:A:185:VAL:HA	1:A:197:GLU:HG3	1.84	0.59
1:D:234:ARG:NH2	7:D:505:HOH:O	2.38	0.56
1:A:309:THR:O	1:A:310:GLN:CB	2.55	0.54
1:C:223:ASN:OD1	1:C:225:GLU:HG2	2.09	0.53
1:A:157:ASN:OD1	1:A:163:ASN:OD1	2.28	0.52
1:D:40:ARG:NH1	7:D:509:HOH:O	2.43	0.52
1:C:167:GLN:HE22	1:C:216:THR:HA	1.74	0.51
1:A:29:VAL:HG22	1:A:38:SER:O	2.11	0.50
1:C:9:MET:HE1	1:C:59:LEU:HD12	1.92	0.50
1:B:223:ASN:OD1	1:B:225:GLU:HG2	2.12	0.50
1:C:40:ARG:NH1	7:C:506:HOH:O	2.45	0.50
1:B:167:GLN:HE22	1:B:216:THR:HA	1.81	0.45
1:C:9:MET:HE1	1:C:59:LEU:CD1	2.48	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:SER:OG	1:D:128:VAL:HG22	2.17	0.44
1:B:9:MET:HG3	1:B:62:PHE:CE1	2.52	0.44
1:C:167:GLN:HG2	7:C:594:HOH:O	2.18	0.43
1:C:299:PRO:N	1:C:300:PRO:CD	2.82	0.43
1:D:42:HIS:HE1	1:D:90:THR:O	2.02	0.43
1:C:97:GLU:OE2	1:C:101[B]:THR:HG21	2.19	0.43
1:B:189:ARG:HD2	1:B:197:GLU:OE2	2.19	0.42
1:A:27:THR:HG22	1:A:28:GLY:N	2.35	0.42
1:B:30[A]:ARG:NH2	7:B:514:HOH:O	2.52	0.42
1:B:168:GLN:OE1	1:B:210:LYS:HE2	2.20	0.42
1:A:274:SER:OG	1:A:304:GLN:HB2	2.21	0.41
1:A:29:VAL:HG22	1:A:38:SER:C	2.41	0.41
1:B:97:GLU:OE2	1:B:101[B]:THR:HG21	2.21	0.41
1:B:239:SER:O	1:B:243:GLU:HG3	2.21	0.41
6:B:405:MXE:H32	7:B:656:HOH:O	2.21	0.41
1:C:261:LEU:HD21	1:C:270[B]:VAL:HG23	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/323 (91%)	289 (99%)	4 (1%)	0	100	100
1	B	304/323 (94%)	299 (98%)	5 (2%)	0	100	100
1	C	301/323 (93%)	297 (99%)	4 (1%)	0	100	100
1	D	291/323 (90%)	286 (98%)	5 (2%)	0	100	100
All	All	1189/1292 (92%)	1171 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/288 (84%)	238 (98%)	4 (2%)	56	35
1	B	263/288 (91%)	260 (99%)	3 (1%)	70	54
1	C	254/288 (88%)	251 (99%)	3 (1%)	67	50
1	D	233/288 (81%)	231 (99%)	2 (1%)	75	63
All	All	992/1152 (86%)	980 (99%)	12 (1%)	67	50

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	LYS
1	A	177	ILE
1	A	197	GLU
1	A	241	PHE
1	B	177	ILE
1	B	241	PHE
1	B	313	LEU
1	C	177	ILE
1	C	183	SER
1	C	241	PHE
1	D	177	ILE
1	D	241	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	OCS	C	305	1	7,8,9	1.01	0	6,11,13	1.55	2 (33%)
1	OCS	A	305	1	7,8,9	1.02	0	6,11,13	1.96	3 (50%)
1	OCS	D	305	1	7,8,9	0.94	0	6,11,13	1.74	2 (33%)
1	OCS	B	305	1	7,8,9	1.02	0	6,11,13	1.83	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	C	305	1	-	0/4/7/9	-
1	OCS	A	305	1	-	0/4/7/9	-
1	OCS	D	305	1	-	0/4/7/9	-
1	OCS	B	305	1	-	0/4/7/9	-

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	305	OCS	OD3-SG-CB	3.46	111.06	106.94
1	A	305	OCS	OD1-SG-CB	2.87	110.35	106.94
1	A	305	OCS	OD3-SG-CB	2.62	110.05	106.94
1	D	305	OCS	OD2-SG-CB	2.51	109.74	105.74
1	D	305	OCS	OD3-SG-CB	2.39	109.78	106.94
1	C	305	OCS	OD1-SG-CB	2.23	109.59	106.94
1	C	305	OCS	OD3-SG-CB	2.15	109.50	106.94
1	A	305	OCS	OD2-SG-CB	2.11	109.10	105.74

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	OXY	C	402	-	1,1,1	0.59	0	-		
2	AZA	A	401	-	9,12,12	1.22	0	4,17,17	3.81	3 (75%)
2	AZA	B	401	-	9,12,12	1.13	0	4,17,17	4.47	3 (75%)
2	AZA	C	401	-	9,12,12	1.14	0	4,17,17	5.24	3 (75%)
4	SO4	A	403	-	4,4,4	0.15	0	6,6,6	0.23	0
3	OXY	B	402	-	1,1,1	0.71	0	-		
4	SO4	B	403	-	4,4,4	0.16	0	6,6,6	0.37	0
6	MXE	C	404	-	4,4,4	0.46	0	3,3,3	0.44	0
2	AZA	D	401	-	9,12,12	1.05	0	4,17,17	4.35	3 (75%)
6	MXE	B	405	-	4,4,4	0.37	0	3,3,3	0.36	0
3	OXY	D	402	-	1,1,1	0.34	0	-		
3	OXY	A	402	-	1,1,1	0.34	0	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AZA	A	401	-	-	-	0/2/2/2
2	AZA	B	401	-	-	-	0/2/2/2
2	AZA	C	401	-	-	-	0/2/2/2
6	MXE	C	404	-	-	1/2/2/2	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AZA	D	401	-	-	-	0/2/2/2
6	MXE	B	405	-	-	1/2/2/2	-

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	AZA	C2-N1-C6	9.30	122.99	115.14
2	B	401	AZA	C2-N1-C6	7.63	121.59	115.14
2	D	401	AZA	C2-N1-C6	7.25	121.26	115.14
2	A	401	AZA	C2-N1-C6	6.49	120.62	115.14
2	C	401	AZA	C5-C6-N1	-3.80	118.23	123.43
2	B	401	AZA	C5-C6-N1	-3.52	118.62	123.43
2	D	401	AZA	C5-C6-N1	-3.41	118.77	123.43
2	D	401	AZA	N9-N8-N7	-3.38	106.86	111.25
2	A	401	AZA	C5-C6-N1	-2.99	119.34	123.43
2	B	401	AZA	N9-N8-N7	-2.96	107.41	111.25
2	A	401	AZA	N9-N8-N7	-2.52	107.98	111.25
2	C	401	AZA	N9-N8-N7	-2.36	108.18	111.25

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	405	MXE	C1-C2-O2-C3
6	C	404	MXE	C1-C2-O2-C3

There are no ring outliers.



5 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	AZA	1	0
2	B	401	AZA	1	0
2	C	401	AZA	1	0
2	D	401	AZA	1	0
6	B	405	MXE	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is






within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	297/323 (91%)	-0.11	11 (3%)	45	48	14, 24, 45, 84	0
1	B	301/323 (93%)	-0.25	11 (3%)	45	48	8, 19, 41, 78	6 (1%)
1	C	298/323 (92%)	-0.30	11 (3%)	45	48	7, 19, 40, 61	7 (2%)
1	D	295/323 (91%)	-0.04	12 (4%)	42	45	14, 25, 46, 67	0
All	All	1191/1292 (92%)	-0.18	45 (3%)	44	47	7, 22, 43, 84	13 (1%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	291	GLY	8.0
1	A	291	GLY	5.7
1	D	8	VAL	5.5
1	A	8	VAL	5.2
1	B	286	ILE	4.7
1	B	283	VAL	4.7
1	C	283	VAL	4.4
1	D	222	THR	4.4
1	A	310	GLN	4.2
1	C	314	PRO	3.8
1	B	313	LEU	3.8
1	A	283	VAL	3.7
1	D	284	GLU	3.3
1	D	283	VAL	3.3
1	A	284	GLU	3.1
1	A	131	GLY	3.1
1	C	10	TYR	3.0
1	B	285	GLU	3.0
1	D	244	THR	3.0
1	A	132	ASN	3.0
1	B	129	LYS	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	59	LEU	2.9
1	A	130	ASN	2.9
1	D	131	GLY	2.8
1	C	282	ILE	2.7
1	C	8	VAL	2.7
1	D	310	GLN	2.7
1	D	292	LYS	2.6
1	D	134	ALA	2.5
1	A	244	THR	2.5
1	B	10	TYR	2.5
1	D	126	PHE	2.4
1	C	281	LYS	2.3
1	B	282	ILE	2.3
1	D	29	VAL	2.3
1	C	313	LEU	2.3
1	B	130	ASN	2.2
1	C	126	PHE	2.2
1	A	223	ASN	2.2
1	C	129	LYS	2.1
1	C	292	LYS	2.1
1	B	134	ALA	2.1
1	A	151	TYR	2.1
1	D	128	VAL	2.1
1	C	160	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	OCS	B	305	9/10	0.92	0.12	14,24,35,37	0
1	OCS	D	305	9/10	0.92	0.11	20,30,42,49	0
1	OCS	C	305	9/10	0.93	0.13	14,23,36,39	0
1	OCS	A	305	9/10	0.94	0.10	17,28,44,48	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	MXE	C	404	5/5	0.80	0.20	34,44,54,61	0
4	SO4	B	403	5/5	0.84	0.17	42,44,63,66	0
6	MXE	B	405	5/5	0.87	0.17	39,47,49,52	0
4	SO4	A	403	5/5	0.91	0.12	42,46,56,61	0
3	OXY	A	402	2/2	0.95	0.07	13,13,13,21	0
2	AZA	D	401	11/11	0.96	0.05	16,19,21,23	0
2	AZA	C	401	11/11	0.96	0.06	14,15,18,19	0
2	AZA	A	401	11/11	0.97	0.05	16,17,20,21	0
3	OXY	B	402	2/2	0.97	0.05	12,12,12,13	0
3	OXY	C	402	2/2	0.97	0.08	13,13,13,19	0
3	OXY	D	402	2/2	0.97	0.06	13,13,13,23	0
2	AZA	B	401	11/11	0.98	0.04	12,14,17,17	0
5	K	B	404	1/1	0.99	0.23	28,28,28,28	0
5	K	C	403	1/1	0.99	0.20	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around OXY A 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AZA D 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AZA C 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around AZA A 401:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around OXY B 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around OXY C 402:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around OXY D 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.