wwPDB X-ray Structure Validation Summary Report (i)

Oct 5, 2024 – 12:28 PM EDT

PDB ID	:	4YLN
Title	:	E. coli Transcription Initiation Complex - 17-bp spacer and 4-nt RNA
Authors	:	Zuo, Y.; Steitz, T.A.
Deposited on	:	2015-03-05
Resolution	:	5.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 5.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1029 (7.00-4.00)
Clashscore	180529	1069 (7.00-4.00)
Ramachandran outliers	177936	1010 (7.04-3.96)
Sidechain outliers	177891	1004 (7.04-3.94)
RSRZ outliers	164620	1023 (7.00-4.00)
RNA backbone	3690	1172 (7.80-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	242	38% 51% 5% 5%
1	B	242	40% 47% 7% 6%
1	G	242	48% 41% 6% 5%
1	H	242	48% 38% 7% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	M	242	49% 43% 5%
1	N	242	48% 40% 6% 6%
2	C	1342	46% 48% 6%
2	I	1342	48% 47% 5%
2	O	1342	50% 45%
3	D	1407	44% 47% 6%
3	J	1407	41% 48% 7%
3	P	1407	45% 45% 7%
4	E	90	48% 50%
4	K	90	53% 42%
4	Q	90	67% 29%
5	F	628	43% 32% 21%
5	L	628	43% 31% 5% 21%
5	R	628	41% 32% 6% 21%
6	1	49	29% 71%
6	4	49	31% 69%
6	7	49	31% 69%
7	2	49	37% 63%
7	5	49	41% 59%
7	8	49	35% 65%
8	3	4	25% 50% 25%
8	6	4	50% 50%
8	9	4	75% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
9	ZN	J	1502	-	-	X	-

2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 94608 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	0	0
			1787	1112	317	352	6			
1	B	228	Total	C	N	O	S	0	0	0
			1767	1100	312	349	6			
1	G	230	Total	C	N	O	S	0	0	0
			1787	1112	317	352	6			
1	H	228	Total	C	N	O	S	0	0	0
			1767	1100	312	349	6			
1	M	230	Total	C	N	O	S	0	0	0
			1787	1112	317	352	6			
1	N	228	Total	C	N	O	S	0	0	0
			1767	1100	312	349	6			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	ALA	-	expression tag	UNP A7ZSI4
A	-5	HIS	-	expression tag	UNP A7ZSI4
A	-4	HIS	-	expression tag	UNP A7ZSI4
A	-3	HIS	-	expression tag	UNP A7ZSI4
A	-2	HIS	-	expression tag	UNP A7ZSI4
A	-1	HIS	-	expression tag	UNP A7ZSI4
A	0	HIS	-	expression tag	UNP A7ZSI4
B	-6	ALA	-	expression tag	UNP A7ZSI4
B	-5	HIS	-	expression tag	UNP A7ZSI4
B	-4	HIS	-	expression tag	UNP A7ZSI4
B	-3	HIS	-	expression tag	UNP A7ZSI4
B	-2	HIS	-	expression tag	UNP A7ZSI4
B	-1	HIS	-	expression tag	UNP A7ZSI4
B	0	HIS	-	expression tag	UNP A7ZSI4
G	-6	ALA	-	expression tag	UNP A7ZSI4
G	-5	HIS	-	expression tag	UNP A7ZSI4
G	-4	HIS	-	expression tag	UNP A7ZSI4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	-3	HIS	-	expression tag	UNP A7ZSI4
G	-2	HIS	-	expression tag	UNP A7ZSI4
G	-1	HIS	-	expression tag	UNP A7ZSI4
G	0	HIS	-	expression tag	UNP A7ZSI4
H	-6	ALA	-	expression tag	UNP A7ZSI4
H	-5	HIS	-	expression tag	UNP A7ZSI4
H	-4	HIS	-	expression tag	UNP A7ZSI4
H	-3	HIS	-	expression tag	UNP A7ZSI4
H	-2	HIS	-	expression tag	UNP A7ZSI4
H	-1	HIS	-	expression tag	UNP A7ZSI4
H	0	HIS	-	expression tag	UNP A7ZSI4
M	-6	ALA	-	expression tag	UNP A7ZSI4
M	-5	HIS	-	expression tag	UNP A7ZSI4
M	-4	HIS	-	expression tag	UNP A7ZSI4
M	-3	HIS	-	expression tag	UNP A7ZSI4
M	-2	HIS	-	expression tag	UNP A7ZSI4
M	-1	HIS	-	expression tag	UNP A7ZSI4
M	0	HIS	-	expression tag	UNP A7ZSI4
N	-6	ALA	-	expression tag	UNP A7ZSI4
N	-5	HIS	-	expression tag	UNP A7ZSI4
N	-4	HIS	-	expression tag	UNP A7ZSI4
N	-3	HIS	-	expression tag	UNP A7ZSI4
N	-2	HIS	-	expression tag	UNP A7ZSI4
N	-1	HIS	-	expression tag	UNP A7ZSI4
N	0	HIS	-	expression tag	UNP A7ZSI4

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1341	Total	C	N	O	S	0	0	0
			10576	6636	1842	2055	43			
2	I	1341	Total	C	N	O	S	0	0	0
			10576	6636	1842	2055	43			
2	O	1341	Total	C	N	O	S	0	0	0
			10576	6636	1842	2055	43			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1362	Total	C	N	O	S	0	0	0
			10568	6633	1887	1998	50			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	J	1362	Total	C	N	O	S	0	0	0
			10568	6633	1887	1998	50			
3	P	1362	Total	C	N	O	S	0	0	0
			10568	6633	1887	1998	50			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	90	Total	C	N	O	S	0	0	0
			708	430	136	141	1			
4	K	90	Total	C	N	O	S	0	0	0
			708	430	136	141	1			
4	Q	90	Total	C	N	O	S	0	0	0
			708	430	136	141	1			

- Molecule 5 is a protein called RNA polymerase sigma factor RpoD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	497	Total	C	N	O	S	0	0	0
			4022	2512	719	768	23			
5	L	497	Total	C	N	O	S	0	0	0
			4022	2512	719	768	23			
5	R	497	Total	C	N	O	S	0	0	0
			4022	2512	719	768	23			

There are 45 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-14	MET	-	expression tag	UNP P00579
F	-13	ARG	-	expression tag	UNP P00579
F	-12	GLY	-	expression tag	UNP P00579
F	-11	SER	-	expression tag	UNP P00579
F	-10	HIS	-	expression tag	UNP P00579
F	-9	HIS	-	expression tag	UNP P00579
F	-8	HIS	-	expression tag	UNP P00579
F	-7	HIS	-	expression tag	UNP P00579
F	-6	HIS	-	expression tag	UNP P00579
F	-5	HIS	-	expression tag	UNP P00579
F	-4	THR	-	expression tag	UNP P00579
F	-3	ASP	-	expression tag	UNP P00579
F	-2	GLN	-	expression tag	UNP P00579
F	-1	PHE	-	expression tag	UNP P00579

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	0	THR	-	expression tag	UNP P00579
L	-14	MET	-	expression tag	UNP P00579
L	-13	ARG	-	expression tag	UNP P00579
L	-12	GLY	-	expression tag	UNP P00579
L	-11	SER	-	expression tag	UNP P00579
L	-10	HIS	-	expression tag	UNP P00579
L	-9	HIS	-	expression tag	UNP P00579
L	-8	HIS	-	expression tag	UNP P00579
L	-7	HIS	-	expression tag	UNP P00579
L	-6	HIS	-	expression tag	UNP P00579
L	-5	HIS	-	expression tag	UNP P00579
L	-4	THR	-	expression tag	UNP P00579
L	-3	ASP	-	expression tag	UNP P00579
L	-2	GLN	-	expression tag	UNP P00579
L	-1	PHE	-	expression tag	UNP P00579
L	0	THR	-	expression tag	UNP P00579
R	-14	MET	-	expression tag	UNP P00579
R	-13	ARG	-	expression tag	UNP P00579
R	-12	GLY	-	expression tag	UNP P00579
R	-11	SER	-	expression tag	UNP P00579
R	-10	HIS	-	expression tag	UNP P00579
R	-9	HIS	-	expression tag	UNP P00579
R	-8	HIS	-	expression tag	UNP P00579
R	-7	HIS	-	expression tag	UNP P00579
R	-6	HIS	-	expression tag	UNP P00579
R	-5	HIS	-	expression tag	UNP P00579
R	-4	THR	-	expression tag	UNP P00579
R	-3	ASP	-	expression tag	UNP P00579
R	-2	GLN	-	expression tag	UNP P00579
R	-1	PHE	-	expression tag	UNP P00579
R	0	THR	-	expression tag	UNP P00579

- Molecule 6 is a DNA chain called NT strand DNA (49-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	1	49	Total	C	N	O	P	0	0	0
			996	476	178	294	48			
6	4	49	Total	C	N	O	P	0	0	0
			996	476	178	294	48			
6	7	49	Total	C	N	O	P	0	0	0
			996	476	178	294	48			

- Molecule 7 is a DNA chain called T strand DNA (49-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	2	49	Total	C	N	O	P	0	0	0
			1012	481	191	292	48			
7	5	49	Total	C	N	O	P	0	0	0
			1012	481	191	292	48			
7	8	49	Total	C	N	O	P	0	0	0
			1012	481	191	292	48			

- Molecule 8 is a RNA chain called RNA (5'-D(*(GTP))-R(P*AP*GP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	3	4	Total	C	N	O	P	0	0	0
			97	39	17	35	6			
8	6	4	Total	C	N	O	P	0	0	0
			97	39	17	35	6			
8	9	4	Total	C	N	O	P	0	0	0
			97	39	17	35	6			

- Molecule 9 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	D	2	Total	Zn	0	0
			2	2		
9	J	2	Total	Zn	0	0
			2	2		
9	P	2	Total	Zn	0	0
			2	2		

- Molecule 10 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	D	1	Total	Mg	0	0
			1	1		
10	6	1	Total	Mg	0	0
			1	1		
10	P	1	Total	Mg	0	0
			1	1		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA-directed RNA polymerase subunit alpha

Chain A:



• Molecule 1: DNA-directed RNA polymerase subunit alpha

Chain B:



• Molecule 1: DNA-directed RNA polymerase subunit alpha

Chain G:



• Molecule 1: DNA-directed RNA polymerase subunit alpha

Chain H:



• Molecule 1: DNA-directed RNA polymerase subunit alpha

Chain M:



• Molecule 1: DNA-directed RNA polymerase subunit alpha

Chain N:





• Molecule 2: DNA-directed RNA polymerase subunit beta

Chain C:





• Molecule 2: DNA-directed RNA polymerase subunit beta

Chain I:





• Molecule 2: DNA-directed RNA polymerase subunit beta

Chain O:





• Molecule 3: DNA-directed RNA polymerase subunit beta'

Chain D:





• Molecule 3: DNA-directed RNA polymerase subunit beta'

Chain J:







• Molecule 3: DNA-directed RNA polymerase subunit beta'

Chain P:





• Molecule 4: DNA-directed RNA polymerase subunit omega

Chain E:



• Molecule 4: DNA-directed RNA polymerase subunit omega

Chain K:



• Molecule 4: DNA-directed RNA polymerase subunit omega

Chain Q:



• Molecule 5: RNA polymerase sigma factor RpoD

Chain F:





• Molecule 5: RNA polymerase sigma factor RpoD

Chain L:







• Molecule 5: RNA polymerase sigma factor RpoD

Chain R:



• Molecule 6: NT strand DNA (49-MER)

Chain 1:





• Molecule 6: NT strand DNA (49-MER)

Chain 4:



• Molecule 6: NT strand DNA (49-MER)

Chain 7:



• Molecule 7: T strand DNA (49-MER)

Chain 2:



• Molecule 7: T strand DNA (49-MER)

Chain 5:



• Molecule 7: T strand DNA (49-MER)

Chain 8:



• Molecule 8: RNA (5'-D(*(GTP))-R(P*AP*GP*U)-3')

Chain 3:



• Molecule 8: RNA (5'-D(*(GTP))-R(P*AP*GP*U)-3')

Chain 6:

• Molecule 8: RNA (5'-D(*(GTP))-R(P*AP*GP*U)-3')

Chain 9:

G13
A14
G15
U16

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	237.40Å 206.05Å 248.69Å 90.00° 116.55° 90.00°	Depositor
Resolution (Å)	39.90 – 5.50 39.90 – 5.50	Depositor EDS
% Data completeness (in resolution range)	99.3 (39.90-5.50) 99.3 (39.90-5.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.20 (at 5.37Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.245 , 0.328 0.245 , 0.327	Depositor DCC
R_{free} test set	3459 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å^2)	268.1	Xtriage
Anisotropy	0.597	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.28 , 190.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.40$, $< L^2 > = 0.23$	Xtriage
Estimated twinning fraction	0.045 for h,-k,-h-l	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	94608	wwPDB-VP
Average B, all atoms (Å^2)	219.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.59	0/1809	0.84	1/2450 (0.0%)
1	B	0.54	0/1789	0.78	0/2425
1	G	0.56	0/1809	0.76	1/2450 (0.0%)
1	H	0.53	0/1789	0.76	0/2425
1	M	0.53	0/1809	0.74	0/2450
1	N	0.54	0/1789	0.79	2/2425 (0.1%)
2	C	0.54	0/10745	0.78	4/14499 (0.0%)
2	I	0.54	3/10745 (0.0%)	0.77	2/14499 (0.0%)
2	O	0.53	0/10745	0.75	3/14499 (0.0%)
3	D	0.54	0/10729	0.77	4/14487 (0.0%)
3	J	0.58	2/10729 (0.0%)	0.81	10/14487 (0.1%)
3	P	0.55	1/10729 (0.0%)	0.77	6/14487 (0.0%)
4	E	0.54	1/710 (0.1%)	0.72	0/956
4	K	0.53	0/710	0.73	0/956
4	Q	0.52	0/710	0.72	0/956
5	F	0.49	1/4076 (0.0%)	0.69	0/5482
5	L	0.51	0/4076	0.72	0/5482
5	R	0.55	2/4076 (0.0%)	0.74	1/5482 (0.0%)
6	1	0.41	0/1115	0.69	0/1718
6	4	0.33	0/1112	0.66	0/1706
6	7	0.37	0/1114	0.67	0/1714
7	2	0.37	0/1134	0.67	0/1744
7	5	0.35	0/1134	0.65	0/1744
7	8	0.38	0/1136	0.64	0/1752
8	3	0.44	0/72	0.62	0/110
8	6	0.40	0/72	0.61	0/110
8	9	0.36	0/72	0.59	0/110
All	All	0.53	10/96535 (0.0%)	0.76	34/131605 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected

by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	D	0	1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	1340	LYS	CB-CG	6.65	1.70	1.52
2	I	626	GLU	CD-OE2	6.62	1.32	1.25
2	I	626	GLU	CD-OE1	5.92	1.32	1.25
5	R	109	GLU	CD-OE1	5.75	1.31	1.25
5	F	491	GLU	CB-CG	5.70	1.62	1.52

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	J	737	ILE	CB-CA-C	-7.98	95.64	111.60
3	J	803	VAL	CB-CA-C	-7.38	97.38	111.40
5	R	488	LEU	CA-CB-CG	7.32	132.12	115.30
3	D	737	ILE	CB-CA-C	-7.15	97.30	111.60
2	O	57	PHE	C-N-CD	-7.09	105.00	120.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	D	671	GLY	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1787	0	1813	220	0
1	B	1767	0	1789	175	0
1	G	1787	0	1812	173	0
1	H	1767	0	1789	149	0
1	M	1787	0	1813	178	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	N	1767	0	1789	142	0
2	C	10576	0	10591	868	0
2	I	10576	0	10591	845	0
2	O	10576	0	10591	771	0
3	D	10568	0	10782	856	3
3	J	10568	0	10780	1069	2
3	P	10568	0	10780	901	0
4	E	708	0	719	42	0
4	K	708	0	719	48	0
4	Q	708	0	719	36	0
5	F	4022	0	4083	243	0
5	L	4022	0	4083	270	0
5	R	4022	0	4083	282	0
6	1	996	0	554	70	1
6	4	996	0	557	76	0
6	7	996	0	555	74	0
7	2	1012	0	556	62	0
7	5	1012	0	556	59	0
7	8	1012	0	554	64	0
8	3	97	0	44	7	0
8	6	97	0	44	8	0
8	9	97	0	44	4	0
9	D	2	0	0	0	0
9	J	2	0	0	2	0
9	P	2	0	0	0	0
10	6	1	0	0	0	0
10	D	1	0	0	0	0
10	P	1	0	0	0	0
All	All	94608	0	92790	6821	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 36.

The worst 5 of 6821 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:608:CYS:SG	3:D:617:THR:HG22	1.31	1.67
3:D:501:VAL:CG1	3:D:502:PRO:HD2	1.33	1.55
3:J:349:TYR:O	3:J:470:VAL:HG23	1.24	1.30
3:D:645:VAL:CG2	3:D:701:LEU:HD13	1.59	1.30
5:L:573:LEU:HB2	7:5:46:DG:OP2	1.15	1.28

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:1169:THR:OG1	6:1:16:DA:OP1[2_657]	1.85	0.35
3:D:710:ASP:OD2	3:J:1282:TYR:OH[2_547]	1.93	0.27
3:D:710:ASP:CA	3:J:1302:TYR:OH[2_547]	2.07	0.13

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	228/242 (94%)	214 (94%)	11 (5%)	3 (1%)	10 42
1	B	226/242 (93%)	204 (90%)	17 (8%)	5 (2%)	5 29
1	G	228/242 (94%)	209 (92%)	16 (7%)	3 (1%)	10 42
1	H	226/242 (93%)	207 (92%)	13 (6%)	6 (3%)	4 25
1	M	228/242 (94%)	214 (94%)	14 (6%)	0	100 100
1	N	226/242 (93%)	209 (92%)	14 (6%)	3 (1%)	10 42
2	C	1339/1342 (100%)	1218 (91%)	98 (7%)	23 (2%)	7 36
2	I	1339/1342 (100%)	1214 (91%)	105 (8%)	20 (2%)	8 39
2	O	1339/1342 (100%)	1234 (92%)	90 (7%)	15 (1%)	12 46
3	D	1360/1407 (97%)	1220 (90%)	109 (8%)	31 (2%)	5 28
3	J	1360/1407 (97%)	1227 (90%)	99 (7%)	34 (2%)	4 26
3	P	1360/1407 (97%)	1226 (90%)	99 (7%)	35 (3%)	4 25
4	E	88/90 (98%)	83 (94%)	5 (6%)	0	100 100
4	K	88/90 (98%)	84 (96%)	3 (3%)	1 (1%)	12 46
4	Q	88/90 (98%)	84 (96%)	4 (4%)	0	100 100
5	F	493/628 (78%)	444 (90%)	27 (6%)	22 (4%)	2 17
5	L	493/628 (78%)	447 (91%)	28 (6%)	18 (4%)	2 20

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
5	R	493/628 (78%)	449 (91%)	30 (6%)	14 (3%)	4 24
All	All	11202/11853 (94%)	10187 (91%)	782 (7%)	233 (2%)	5 29

5 of 233 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	164	ASP
1	B	118	ASP
2	C	110	PRO
2	C	214	ASN
2	C	247	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	198/208 (95%)	181 (91%)	17 (9%)	8 26
1	B	196/208 (94%)	172 (88%)	24 (12%)	4 16
1	G	198/208 (95%)	178 (90%)	20 (10%)	6 20
1	H	196/208 (94%)	174 (89%)	22 (11%)	5 17
1	M	198/208 (95%)	178 (90%)	20 (10%)	6 20
1	N	196/208 (94%)	176 (90%)	20 (10%)	6 20
2	C	1156/1157 (100%)	1042 (90%)	114 (10%)	6 21
2	I	1156/1157 (100%)	1052 (91%)	104 (9%)	8 24
2	O	1156/1157 (100%)	1050 (91%)	106 (9%)	7 24
3	D	1135/1168 (97%)	1026 (90%)	109 (10%)	7 22
3	J	1135/1168 (97%)	1014 (89%)	121 (11%)	5 19
3	P	1135/1168 (97%)	1017 (90%)	118 (10%)	5 20
4	E	74/74 (100%)	70 (95%)	4 (5%)	18 40
4	K	74/74 (100%)	67 (90%)	7 (10%)	7 22
4	Q	74/74 (100%)	66 (89%)	8 (11%)	5 19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
5	F	439/554 (79%)	406 (92%)	33 (8%)	11 31
5	L	439/554 (79%)	394 (90%)	45 (10%)	6 20
5	R	439/554 (79%)	393 (90%)	46 (10%)	5 19
All	All	9594/10107 (95%)	8656 (90%)	938 (10%)	6 21

5 of 938 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	J	153	ASN
4	Q	19	LEU
4	K	65	ASP
3	P	1321	SER
3	P	442	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 148 such sidechains are listed below:

Mol	Chain	Res	Type
3	P	113	HIS
5	R	383	ASN
3	P	294	ASN
3	P	736	GLN
1	G	66	HIS

5.3.3 RNA

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
8	3	2/4 (50%)	1 (50%)	0
8	6	3/4 (75%)	1 (33%)	1 (33%)
8	9	3/4 (75%)	1 (33%)	1 (33%)
All	All	8/12 (66%)	3 (37%)	2 (25%)

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
8	3	15	G
8	6	15	G
8	9	15	G

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
8	6	13	GTP
8	9	13	GTP

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
6	4	3
7	2	3
7	5	3
6	7	1
7	8	1

The worst 5 of 11 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	4	45:DT	O3'	46:DG	P	5.04
1	7	50:DT	O3'	51:DC	P	4.24
1	8	22:DA	O3'	23:DT	P	3.80
1	2	22:DA	O3'	23:DT	P	3.79
1	5	22:DA	O3'	23:DT	P	3.79

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	230/242 (95%)	-0.83	0 100 100	153, 175, 210, 235	0
1	B	228/242 (94%)	-0.77	1 (0%) 89 79	162, 194, 217, 238	0
1	G	230/242 (95%)	-0.69	0 100 100	157, 185, 216, 248	0
1	H	228/242 (94%)	-0.73	0 100 100	160, 191, 229, 261	0
1	M	230/242 (95%)	-0.77	0 100 100	166, 200, 233, 252	0
1	N	228/242 (94%)	-0.57	0 100 100	186, 233, 258, 273	0
2	C	1341/1342 (99%)	-0.74	2 (0%) 92 87	119, 186, 244, 277	0
2	I	1341/1342 (99%)	-0.76	0 100 100	130, 195, 278, 377	0
2	O	1341/1342 (99%)	-0.74	1 (0%) 92 87	144, 183, 235, 270	0
3	D	1362/1407 (96%)	-0.67	3 (0%) 92 84	128, 214, 296, 349	0
3	J	1362/1407 (96%)	-0.67	1 (0%) 92 87	132, 194, 280, 314	0
3	P	1362/1407 (96%)	-0.67	3 (0%) 92 84	148, 208, 292, 330	0
4	E	90/90 (100%)	-0.54	0 100 100	169, 206, 407, 461	0
4	K	90/90 (100%)	-0.58	0 100 100	144, 199, 394, 442	0
4	Q	90/90 (100%)	-0.70	0 100 100	167, 222, 416, 460	0
5	F	497/628 (79%)	-0.62	0 100 100	182, 294, 404, 418	0
5	L	497/628 (79%)	-0.59	1 (0%) 92 84	168, 262, 400, 406	0
5	R	497/628 (79%)	-0.61	0 100 100	172, 259, 413, 444	0
6	1	49/49 (100%)	-0.39	0 100 100	201, 272, 311, 317	0
6	4	49/49 (100%)	-0.35	0 100 100	209, 264, 308, 350	0
6	7	49/49 (100%)	-0.27	0 100 100	211, 255, 278, 300	0
7	2	49/49 (100%)	-0.38	0 100 100	210, 278, 312, 343	0
7	5	49/49 (100%)	-0.37	0 100 100	195, 270, 339, 341	0
7	8	49/49 (100%)	-0.35	0 100 100	194, 260, 296, 335	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
8	3	3/4 (75%)	-0.83	0	100	100	255, 255, 281, 321	0
8	6	3/4 (75%)	-0.73	0	100	100	263, 263, 272, 282	0
8	9	3/4 (75%)	-0.59	0	100	100	262, 262, 277, 295	0
All	All	11547/12159 (94%)	-0.69	12 (0%)	92	87	119, 203, 358, 461	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	282	LEU	5.4
3	D	772	TYR	4.8
5	L	304	THR	3.1
3	P	971	GLY	3.0
1	B	90	VAL	2.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
10	MG	6	101	1/1	0.96	0.22	189,189,189,189	0
9	ZN	D	1502	1/1	0.99	0.02	212,212,212,212	0
9	ZN	J	1501	1/1	0.99	0.06	200,200,200,200	0
9	ZN	D	1501	1/1	0.99	0.04	228,228,228,228	0
10	MG	P	1503	1/1	0.99	0.03	194,194,194,194	0
9	ZN	P	1502	1/1	1.00	0.06	187,187,187,187	0
10	MG	D	1503	1/1	1.00	0.02	176,176,176,176	0
9	ZN	J	1502	1/1	1.00	0.06	174,174,174,174	0
9	ZN	P	1501	1/1	1.00	0.02	214,214,214,214	0

6.5 Other polymers (i)

There are no such residues in this entry.