



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 24, 2024 – 10:51 AM EDT

PDB ID : 6Z72  
Title : SARS-CoV-2 Macrodomein in complex with ADP-HPM  
Authors : Zorzini, V.; Rack, J.; Ahel, I.  
Deposited on : 2020-05-29  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	176	<div> <div>%</div> <div> <div></div> <div>89%</div> <div>10%</div> <div>.</div> </div> </div>
1	B	176	<div> <div>2%</div> <div> <div></div> <div>87%</div> <div>11%</div> <div>.</div> </div> </div>
1	C	176	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>9%</div> <div>.</div> </div> </div>
1	D	176	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	403	-	X	-	-
3	EDO	A	404	-	X	-	-
3	EDO	C	402	-	X	-	-
3	EDO	C	403	-	X	-	-
3	EDO	D	403	-	X	-	-
5	MLT	D	402	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5554 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Replicase polypeptide 1ab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	174	Total	C	N	O	S	0	0	0
			1271	805	213	248	5			
1	B	173	Total	C	N	O	S	0	0	0
			1268	804	214	245	5			
1	C	174	Total	C	N	O	S	0	1	1
			1277	810	213	249	5			
1	D	172	Total	C	N	O	S	0	1	0
			1269	807	212	245	5			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	204	GLY	-	expression tag	UNP P0DTD1
A	205	PRO	-	expression tag	UNP P0DTD1
B	204	GLY	-	expression tag	UNP P0DTD1
B	205	PRO	-	expression tag	UNP P0DTD1
C	204	GLY	-	expression tag	UNP P0DTD1
C	205	PRO	-	expression tag	UNP P0DTD1
D	204	GLY	-	expression tag	UNP P0DTD1
D	205	PRO	-	expression tag	UNP P0DTD1

- Molecule 2 is Adenosine Diphosphate (Hydroxymethyl)pyrrolidine monoalcohol (three-letter code: A3R) (formula: C<sub>15</sub>H<sub>24</sub>N<sub>6</sub>O<sub>11</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 34	C 15	N 6	O 11	P 2	0	0
2	B	1	Total 34	C 15	N 6	O 11	P 2	0	0
2	C	1	Total 34	C 15	N 6	O 11	P 2	0	0
2	D	1	Total 34	C 15	N 6	O 11	P 2	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $\text{C}_2\text{H}_6\text{O}_2$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Na 1 1	0	0
4	B	1	Total Na 1 1	0	0
4	C	1	Total Na 1 1	0	0
4	D	1	Total Na 1 1	0	0

- Molecule 5 is D-MALATE (three-letter code: MLT) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			9	4	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	58	Total	O	0	0
			58	58		
6	B	65	Total	O	0	0
			65	65		
6	C	79	Total	O	0	0
			79	79		
6	D	90	Total	O	0	0
			90	90		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Replicase polyprotein 1ab





- Molecule 1: Replicase polyprotein 1ab



- Molecule 1: Replicase polyprotein 1ab



- Molecule 1: Replicase polyprotein 1ab





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.81Å 83.13Å 84.28Å 90.00° 94.41° 90.00°	Depositor
Resolution (Å)	50.50 – 2.30 50.50 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.0 (50.50-2.30) 99.6 (50.50-2.30)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.80 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, $R_{free}$	0.218 , 0.263 0.227 , 0.268	Depositor DCC
$R_{free}$ test set	1834 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.9	Xtriage
Anisotropy	0.792	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 55.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5554	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.87 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0755e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A3R, EDO, NA, MLT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/1293	0.61	0/1765
1	B	0.45	0/1290	0.63	0/1760
1	C	0.49	0/1302	0.66	0/1774
1	D	0.55	0/1294	0.67	0/1764
All	All	0.48	0/5179	0.64	0/7063

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1271	0	1208	15	0
1	B	1268	0	1223	12	0
1	C	1277	0	1227	12	0
1	D	1269	0	1229	13	0
2	A	34	0	0	0	0
2	B	34	0	0	1	0
2	C	34	0	0	1	0
2	D	34	0	0	0	0
3	A	12	0	18	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	4	0	6	0	0
3	C	8	0	12	3	0
3	D	4	0	6	3	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	D	9	0	4	1	0
6	A	58	0	0	1	0
6	B	65	0	0	2	0
6	C	79	0	0	1	0
6	D	90	0	0	1	0
All	All	5554	0	4933	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:217:THR:HG22	1:A:218:ASP:H	1.60	0.67
1:D:306:LYS:NZ	6:D:504:HOH:O	2.29	0.66
1:A:322:GLN:HG3	1:A:323:HIS:CE1	2.36	0.60
1:A:363:ASN:ND2	6:A:503:HOH:O	2.35	0.60
1:A:223:LYS:HG3	3:A:402:EDO:H21	1.83	0.59

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	172/176 (98%)	170 (99%)	2 (1%)	0	100	100
1	B	171/176 (97%)	167 (98%)	4 (2%)	0	100	100
1	C	173/176 (98%)	169 (98%)	4 (2%)	0	100	100
1	D	171/176 (97%)	168 (98%)	3 (2%)	0	100	100
All	All	687/704 (98%)	674 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/147 (89%)	131 (100%)	0	100	100
1	B	133/147 (90%)	133 (100%)	0	100	100
1	C	133/147 (90%)	133 (100%)	0	100	100
1	D	133/147 (90%)	133 (100%)	0	100	100
All	All	530/588 (90%)	530 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	224	ASN
1	C	342	HIS
1	D	323	HIS
1	A	322	GLN
1	A	224	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	EDO	A	402	-	3,3,3	0.58	0	2,2,2	0.72	0
2	A3R	B	401	-	33,37,37	1.05	3 (9%)	35,56,56	1.36	4 (11%)
2	A3R	C	401	-	33,37,37	1.11	2 (6%)	35,56,56	1.34	6 (17%)
3	EDO	C	402	-	3,3,3	2.46	2 (66%)	2,2,2	1.61	0
5	MLT	D	402	-	8,8,8	1.30	1 (12%)	10,10,10	1.76	3 (30%)
3	EDO	A	403	-	3,3,3	2.02	2 (66%)	2,2,2	1.05	0
3	EDO	C	403	-	3,3,3	2.12	2 (66%)	2,2,2	1.05	0
2	A3R	D	401	-	33,37,37	1.07	1 (3%)	35,56,56	1.30	4 (11%)
3	EDO	B	402	-	3,3,3	2.03	2 (66%)	2,2,2	0.81	0
3	EDO	A	404	-	3,3,3	2.13	2 (66%)	2,2,2	1.15	0
3	EDO	D	403	-	3,3,3	2.06	2 (66%)	2,2,2	1.00	0
2	A3R	A	401	-	33,37,37	1.24	1 (3%)	35,56,56	1.32	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	402	-	-	1/1/1/1	-
2	A3R	B	401	-	-	4/18/47/47	0/4/4/4
2	A3R	C	401	-	-	4/18/47/47	0/4/4/4
3	EDO	C	402	-	-	1/1/1/1	-
5	MLT	D	402	-	-	6/8/8/8	-
3	EDO	A	403	-	-	1/1/1/1	-
3	EDO	C	403	-	-	1/1/1/1	-
2	A3R	D	401	-	-	3/18/47/47	0/4/4/4
3	EDO	B	402	-	-	0/1/1/1	-
3	EDO	A	404	-	-	1/1/1/1	-
3	EDO	D	403	-	-	1/1/1/1	-
2	A3R	A	401	-	-	2/18/47/47	0/4/4/4

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	A3R	O4'-C1'	4.70	1.47	1.40
2	D	401	A3R	O4'-C1'	3.79	1.45	1.40
2	C	401	A3R	O4'-C1'	3.62	1.45	1.40
2	B	401	A3R	O4'-C1'	3.35	1.45	1.40
3	C	402	EDO	O2-C2	3.03	1.57	1.42

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	A3R	N3-C2-N1	-5.12	121.73	128.67
2	B	401	A3R	N3-C2-N1	-5.07	121.79	128.67
2	C	401	A3R	N3-C2-N1	-4.34	122.78	128.67
2	D	401	A3R	N3-C2-N1	-4.08	123.14	128.67
5	D	402	MLT	O2-C1-C2	3.95	121.08	112.74

There are no chirality outliers.

5 of 25 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	D	402	MLT	O3-C2-C3-C4
5	D	402	MLT	C1-C2-C3-C4
3	A	402	EDO	O1-C1-C2-O2
3	C	402	EDO	O1-C1-C2-O2
3	C	403	EDO	O1-C1-C2-O2

There are no ring outliers.


8 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	EDO	2	0
2	B	401	A3R	1	0
2	C	401	A3R	1	0
3	C	402	EDO	2	0
5	D	402	MLT	1	0
3	A	403	EDO	2	0
3	C	403	EDO	1	0
3	D	403	EDO	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




## Ligand A3R C 401



## Ligand A3R D 401







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	174/176 (98%)	0.50	1 (0%) 89 92	47, 53, 65, 77	0
1	B	173/176 (98%)	0.48	3 (1%) 70 76	38, 52, 61, 79	0
1	C	174/176 (98%)	0.10	2 (1%) 80 85	21, 32, 51, 88	0
1	D	172/176 (97%)	0.16	2 (1%) 79 83	20, 31, 49, 63	0
All	All	693/704 (98%)	0.31	8 (1%) 79 83	20, 49, 60, 88	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	207	VAL	3.1
1	A	338	ALA	2.8
1	B	345	ARG	2.8
1	C	218	ASP	2.5
1	D	304	VAL	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	MLT	D	402	9/9	0.43	0.42	80,83,88,89	0
3	EDO	C	402	4/4	0.75	0.24	29,41,42,48	0
4	NA	B	403	1/1	0.78	0.18	51,51,51,51	0
3	EDO	D	403	4/4	0.80	0.22	39,41,45,53	0
3	EDO	A	402	4/4	0.83	0.22	51,58,58,62	0
3	EDO	A	404	4/4	0.83	0.22	55,58,60,63	0
4	NA	D	404	1/1	0.86	0.14	33,33,33,33	0
3	EDO	C	403	4/4	0.87	0.18	47,49,53,57	0
3	EDO	A	403	4/4	0.88	0.15	68,70,73,75	0
3	EDO	B	402	4/4	0.89	0.33	62,63,63,65	0
4	NA	A	405	1/1	0.91	0.11	46,46,46,46	0
2	A3R	A	401	34/34	0.95	0.14	47,49,56,64	0
2	A3R	D	401	34/34	0.96	0.14	20,33,40,46	0
2	A3R	B	401	34/34	0.96	0.14	47,47,56,71	0
2	A3R	C	401	34/34	0.97	0.14	18,29,39,41	0
4	NA	C	404	1/1	0.97	0.15	51,51,51,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around A3R A 401:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around A3R D 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around A3R B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.