



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 31, 2025 – 08:56 PM JST

PDB ID : 5Z94 / pdb_00005z94
Title : Crystal Structure of SIRT3 in complex with H3K4bhb peptide
Authors : Zhang, X.; Li, H.
Deposited on : 2018-02-02
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	 92%
1	B	283	 94%
2	C	15	 40% 7% 53%
2	D	15	 40% 7% 53%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-dependent protein deacetylase sirtuin-3, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	273	Total	C	N	O	S	0	1	0
			2154	1388	372	385	9			
1	B	273	Total	C	N	O	S	0	1	0
			2154	1388	372	385	9			

- Molecule 2 is a protein called Gene for histone H3 (germline gene).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	7	Total	C	N	O	0	0	0
			59	35	12	12			
2	D	7	Total	C	N	O	0	0	0
			59	35	12	12			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0

- Molecule 6 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 13 6 7	0	0
6	B	1	Total C O 13 6 7	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	240	Total O 240 240	0	0
7	B	238	Total O 238 238	0	0
7	C	12	Total O 12 12	0	0
7	D	17	Total O 17 17	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial

Chain A:  92%




- Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial

Chain B:  94%




- Molecule 2: Gene for histone H3 (germline gene)

Chain C:  40% 7% 53%



- Molecule 2: Gene for histone H3 (germline gene)

Chain D:  40% 7% 53%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.47Å 76.26Å 75.42Å 90.00° 115.77° 90.00°	Depositor
Resolution (Å)	37.64 – 1.90 37.64 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (37.64-1.90) 99.4 (37.64-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 1.89Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.161 , 0.191 0.162 , 0.191	Depositor DCC
R_{free} test set	2779 reflections (4.59%)	wwPDB-VP
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.460 for l,-k,h	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4995	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, KHB, ZN, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2210	0.54	1/3011 (0.0%)
1	B	0.35	0/2210	0.53	0/3011
2	C	0.26	0/42	0.53	0/54
2	D	0.24	0/42	0.52	0/54
All	All	0.35	0/4504	0.53	1/6130 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	LEU	CA-CB-CG	5.67	128.35	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2154	0	2166	7	0
1	B	2154	0	2166	3	0
2	C	59	0	47	0	0
2	D	59	0	47	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	12	0	16	0	0
4	B	6	0	8	0	0
4	D	6	0	8	0	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	A	13	0	5	0	0
6	B	13	0	5	0	0
7	A	240	0	0	0	0
7	B	238	0	0	0	0
7	C	12	0	0	0	0
7	D	17	0	0	0	0
All	All	4995	0	4468	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:160:PRO:HA	1:B:165:TYR:CG	2.46	0.50
1:A:159:SER:HA	1:A:160:PRO:HD3	1.80	0.45
1:A:150:THR:OG1	1:A:151:PRO:HD3	2.19	0.43
1:A:160:PRO:HA	1:A:165:TYR:CG	2.54	0.43
1:B:150:THR:OG1	1:B:151:PRO:HD3	2.18	0.43
1:A:160:PRO:HA	1:A:165:TYR:CD1	2.56	0.40
1:A:177:GLU:H	1:A:177:GLU:CD	2.24	0.40
1:B:156:ASP:CG	1:B:158:ARG:HH22	2.24	0.40
1:A:216:LEU:HD12	1:A:222:LEU:HB2	2.02	0.40
1:A:344:ASN:O	1:A:363:LEU:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/283 (96%)	268 (98%)	4 (2%)	0	100	100
1	B	272/283 (96%)	269 (99%)	3 (1%)	0	100	100
2	C	4/15 (27%)	4 (100%)	0	0	100	100
2	D	4/15 (27%)	4 (100%)	0	0	100	100
All	All	552/596 (93%)	545 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/245 (97%)	235 (99%)	3 (1%)	65	65
1	B	238/245 (97%)	236 (99%)	2 (1%)	79	80
2	C	4/9 (44%)	4 (100%)	0	100	100
2	D	4/9 (44%)	4 (100%)	0	100	100
All	All	484/508 (95%)	479 (99%)	5 (1%)	73	74

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	158	ARG
1	A	303	LEU
1	A	346	ASP
1	B	346	ASP
1	B	391	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	KHB	C	4	2	13,14,15	1.52	2 (15%)	10,16,18	1.36	2 (20%)
2	KHB	D	4	2	13,14,15	1.54	2 (15%)	10,16,18	1.45	3 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	KHB	C	4	2	-	0/13/14/16	-
2	KHB	D	4	2	-	0/13/14/16	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	4	KHB	C1-NZ	4.49	1.43	1.33
2	C	4	KHB	C1-NZ	4.43	1.43	1.33
2	D	4	KHB	CA-N	-2.27	1.41	1.48
2	C	4	KHB	CA-N	-2.13	1.41	1.48

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4	KHB	CE-NZ-C1	-2.56	118.09	122.84
2	D	4	KHB	C2-C1-NZ	2.26	119.10	115.97
2	D	4	KHB	O1-C1-NZ	-2.26	118.75	123.01

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4	KHB	C2-C1-NZ	2.12	118.91	115.97
2	C	4	KHB	CE-NZ-C1	-2.11	118.93	122.84

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	A	404	-	4,4,4	0.18	0	6,6,6	0.15	0
6	CIT	B	404	-	12,12,12	0.97	0	17,17,17	1.62	2 (11%)
4	GOL	D	101	-	5,5,5	1.00	0	5,5,5	0.83	0
4	GOL	A	403	-	5,5,5	0.97	0	5,5,5	1.15	1 (20%)
6	CIT	A	405	-	12,12,12	0.97	0	17,17,17	1.51	2 (11%)
4	GOL	A	402	-	5,5,5	0.93	0	5,5,5	0.98	0
4	GOL	B	402	-	5,5,5	0.95	0	5,5,5	1.22	1 (20%)
5	SO4	B	403	-	4,4,4	0.20	0	6,6,6	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	CIT	B	404	-	-	2/16/16/16	-
4	GOL	D	101	-	-	1/4/4/4	-
4	GOL	A	403	-	-	4/4/4/4	-
6	CIT	A	405	-	-	8/16/16/16	-
4	GOL	A	402	-	-	0/4/4/4	-
4	GOL	B	402	-	-	4/4/4/4	-

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	404	CIT	O6-C6-C3	4.67	121.15	113.05
6	A	405	CIT	O6-C6-C3	4.12	120.21	113.05
4	B	402	GOL	C3-C2-C1	-2.27	102.87	111.70
6	A	405	CIT	O2-C1-C2	2.21	121.43	114.35
4	A	403	GOL	C3-C2-C1	-2.19	103.20	111.70
6	B	404	CIT	O4-C5-C4	2.13	121.20	114.35

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	403	GOL	O1-C1-C2-C3
4	A	403	GOL	C1-C2-C3-O3
4	B	402	GOL	O1-C1-C2-C3
4	B	402	GOL	C1-C2-C3-O3
4	D	101	GOL	O1-C1-C2-C3
4	A	403	GOL	O1-C1-C2-O2
6	A	405	CIT	O7-C3-C6-O5
6	A	405	CIT	C2-C3-C6-O5
6	A	405	CIT	C2-C3-C6-O6
6	A	405	CIT	C4-C3-C6-O6
6	A	405	CIT	O1-C1-C2-C3
6	A	405	CIT	O2-C1-C2-C3
4	A	403	GOL	O2-C2-C3-O3
6	A	405	CIT	C4-C3-C6-O5
4	B	402	GOL	O1-C1-C2-O2
6	A	405	CIT	O7-C3-C6-O6
4	B	402	GOL	O2-C2-C3-O3
6	B	404	CIT	C2-C3-C6-O6
6	B	404	CIT	C4-C3-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/283 (96%)	-1.28	0 100 100	14, 27, 76, 110	1 (0%)
1	B	273/283 (96%)	-1.29	0 100 100	14, 27, 74, 110	1 (0%)
2	C	6/15 (40%)	-0.31	0 100 100	29, 38, 63, 76	0
2	D	6/15 (40%)	-0.44	0 100 100	29, 39, 67, 75	0
All	All	558/596 (93%)	-1.26	0 100 100	14, 27, 76, 110	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	KHB	C	4	15/16	0.99	0.03	21,23,26,32	0
2	KHB	D	4	15/16	0.99	0.03	20,23,30,33	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	A	402	6/6	0.97	0.05	40,54,60,66	0
4	GOL	B	402	6/6	0.97	0.06	38,41,50,51	0
4	GOL	D	101	6/6	0.97	0.06	40,54,61,63	0
6	CIT	B	404	13/13	0.97	0.06	67,78,88,91	0
6	CIT	A	405	13/13	0.98	0.05	64,78,89,91	0
4	GOL	A	403	6/6	0.98	0.04	34,40,47,49	0
5	SO4	A	404	5/5	0.99	0.03	45,47,48,59	0
5	SO4	B	403	5/5	0.99	0.03	46,48,50,57	0
3	ZN	A	401	1/1	1.00	0.01	20,20,20,20	0
3	ZN	B	401	1/1	1.00	0.01	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.