



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 26, 2024 – 02:57 AM EDT

PDB ID : 6ZDX
Title : RIFIN variable region bound to LILRB1 ectodomain
Authors : Harrison, T.E.; Higgins, M.K.
Deposited on : 2020-06-15
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	129	
2	B	437	
3	C	2	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rifin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	0	0
			777	499	128	144	6			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	145	GLY	-	expression tag	UNP Q8I4N9
A	146	SER	-	expression tag	UNP Q8I4N9
A	147	SER	-	expression tag	UNP Q8I4N9
A	148	HIS	-	expression tag	UNP Q8I4N9
A	149	HIS	-	expression tag	UNP Q8I4N9
A	150	HIS	-	expression tag	UNP Q8I4N9
A	151	HIS	-	expression tag	UNP Q8I4N9
A	152	HIS	-	expression tag	UNP Q8I4N9
A	153	HIS	-	expression tag	UNP Q8I4N9
A	154	SER	-	expression tag	UNP Q8I4N9
A	155	SER	-	expression tag	UNP Q8I4N9
A	156	GLY	-	expression tag	UNP Q8I4N9
A	157	LEU	-	expression tag	UNP Q8I4N9
A	158	VAL	-	expression tag	UNP Q8I4N9
A	159	PRO	-	expression tag	UNP Q8I4N9
A	160	ARG	-	expression tag	UNP Q8I4N9
A	161	GLY	-	expression tag	UNP Q8I4N9
A	162	SER	-	expression tag	UNP Q8I4N9
A	163	HIS	-	expression tag	UNP Q8I4N9
A	164	MET	-	expression tag	UNP Q8I4N9

- Molecule 2 is a protein called Leukocyte immunoglobulin-like receptor subfamily B member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	396	Total	C	N	O	S	0	0	0
			3072	1941	527	592	12			

There are 41 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-30	MET	-	initiating methionine	UNP A0A0G2JQ46
B	-29	GLY	-	expression tag	UNP A0A0G2JQ46
B	-28	ILE	-	expression tag	UNP A0A0G2JQ46
B	-27	LEU	-	expression tag	UNP A0A0G2JQ46
B	-26	PRO	-	expression tag	UNP A0A0G2JQ46
B	-25	SER	-	expression tag	UNP A0A0G2JQ46
B	-24	PRO	-	expression tag	UNP A0A0G2JQ46
B	-23	GLY	-	expression tag	UNP A0A0G2JQ46
B	-22	MET	-	expression tag	UNP A0A0G2JQ46
B	-21	PRO	-	expression tag	UNP A0A0G2JQ46
B	-20	ALA	-	expression tag	UNP A0A0G2JQ46
B	-19	LEU	-	expression tag	UNP A0A0G2JQ46
B	-18	LEU	-	expression tag	UNP A0A0G2JQ46
B	-17	SER	-	expression tag	UNP A0A0G2JQ46
B	-16	LEU	-	expression tag	UNP A0A0G2JQ46
B	-15	VAL	-	expression tag	UNP A0A0G2JQ46
B	-14	SER	-	expression tag	UNP A0A0G2JQ46
B	-13	LEU	-	expression tag	UNP A0A0G2JQ46
B	-12	LEU	-	expression tag	UNP A0A0G2JQ46
B	-11	SER	-	expression tag	UNP A0A0G2JQ46
B	-10	VAL	-	expression tag	UNP A0A0G2JQ46
B	-9	LEU	-	expression tag	UNP A0A0G2JQ46
B	-8	LEU	-	expression tag	UNP A0A0G2JQ46
B	-7	MET	-	expression tag	UNP A0A0G2JQ46
B	-6	GLY	-	expression tag	UNP A0A0G2JQ46
B	-5	CYS	-	expression tag	UNP A0A0G2JQ46
B	-4	VAL	-	expression tag	UNP A0A0G2JQ46
B	-3	ALA	-	expression tag	UNP A0A0G2JQ46
B	-2	GLU	-	expression tag	UNP A0A0G2JQ46
B	-1	THR	-	expression tag	UNP A0A0G2JQ46
B	0	GLY	-	expression tag	UNP A0A0G2JQ46
B	1	GLY	-	expression tag	UNP A0A0G2JQ46
B	398	GLY	-	expression tag	UNP A0A0G2JQ46
B	399	THR	-	expression tag	UNP A0A0G2JQ46
B	400	LYS	-	expression tag	UNP A0A0G2JQ46
B	401	HIS	-	expression tag	UNP A0A0G2JQ46
B	402	HIS	-	expression tag	UNP A0A0G2JQ46
B	403	HIS	-	expression tag	UNP A0A0G2JQ46
B	404	HIS	-	expression tag	UNP A0A0G2JQ46
B	405	HIS	-	expression tag	UNP A0A0G2JQ46
B	406	HIS	-	expression tag	UNP A0A0G2JQ46

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Rifin



- Molecule 2: Leukocyte immunoglobulin-like receptor subfamily B member 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	134.51Å 134.51Å 277.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	89.21 – 3.00 89.21 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (89.21-3.00) 99.9 (89.21-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.234 , 0.249 0.233 , 0.241	Depositor DCC
R_{free} test set	1497 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	97.4	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 88.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3905	wwPDB-VP
Average B, all atoms (Å ²)	120.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/790	0.73	0/1069
2	B	0.51	0/3160	0.74	0/4311
All	All	0.51	0/3950	0.74	0/5380

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	777	0	810	8	0
2	B	3072	0	2951	67	0
3	C	28	0	25	0	0
4	B	28	0	26	0	0
All	All	3905	0	3812	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:308:GLN:HB3	2:B:309:PRO:HD3	1.06	1.06
2:B:308:GLN:HB3	2:B:309:PRO:CD	1.87	1.05
2:B:308:GLN:CB	2:B:309:PRO:HD3	1.97	0.95
2:B:36:ARG:HH11	2:B:36:ARG:HG3	1.43	0.81
2:B:264:VAL:HG13	2:B:268:TYR:CG	2.27	0.69
2:B:366:SER:HA	2:B:396:VAL:HG11	1.74	0.69
2:B:245:LEU:HD22	2:B:259:PHE:HB2	1.77	0.66
2:B:219:LEU:CD2	2:B:264:VAL:HG21	2.27	0.65
2:B:381:LYS:HB3	2:B:384:LEU:HD22	1.79	0.64
2:B:168:ARG:HH11	2:B:170:TRP:HA	1.64	0.63
2:B:32:THR:O	2:B:33:GLN:HB2	1.98	0.62
2:B:219:LEU:HD23	2:B:264:VAL:HG21	1.80	0.62
2:B:219:LEU:HD23	2:B:264:VAL:CG2	2.30	0.61
2:B:264:VAL:HG13	2:B:268:TYR:HB2	1.81	0.61
2:B:119:ILE:HG23	2:B:158:ILE:HG23	1.84	0.60
2:B:36:ARG:HG3	2:B:36:ARG:NH1	2.17	0.59
2:B:328:TRP:CZ2	2:B:330:GLN:HG3	2.38	0.58
2:B:264:VAL:HG22	2:B:268:TYR:CD2	2.39	0.58
2:B:264:VAL:HG13	2:B:268:TYR:CB	2.35	0.57
2:B:138:GLU:HG3	2:B:139:ASP:N	2.19	0.56
2:B:335:THR:HG22	2:B:343:PRO:HB3	1.86	0.55
2:B:303:VAL:HG12	2:B:325:SER:HB3	1.88	0.55
1:A:165:ASN:O	1:A:169:ILE:HG13	2.06	0.55
2:B:207:VAL:HG21	2:B:293:ILE:HD11	1.89	0.55
2:B:337:GLU:HG3	2:B:371:THR:HB	1.87	0.54
2:B:148:GLN:H	2:B:149:PRO:CD	2.21	0.54
2:B:364:VAL:HG12	2:B:396:VAL:HG21	1.90	0.54
2:B:312:THR:HG23	2:B:312:THR:O	2.08	0.53
2:B:202:LYS:NZ	2:B:289:ASP:OD1	2.37	0.53
2:B:78:GLY:HA2	2:B:84:ARG:HA	1.91	0.53
2:B:298:GLN:HG3	2:B:384:LEU:HG	1.91	0.53
2:B:219:LEU:CD2	2:B:264:VAL:CG2	2.88	0.51
2:B:346:LEU:HD22	2:B:359:PHE:CD2	2.46	0.51
2:B:212:ILE:HD12	2:B:335:THR:HG21	1.93	0.51
2:B:65:ILE:HA	2:B:69:HIS:HD2	1.76	0.50
2:B:148:GLN:H	2:B:149:PRO:HD3	1.76	0.50
2:B:76:TYR:CD2	2:B:76:TYR:C	2.85	0.50
2:B:165:SER:H	2:B:170:TRP:HZ2	1.60	0.50
2:B:264:VAL:HG22	2:B:268:TYR:CE2	2.47	0.49
2:B:308:GLN:CB	2:B:309:PRO:CD	2.63	0.49
2:B:138:GLU:CG	2:B:139:ASP:N	2.76	0.49
2:B:148:GLN:N	2:B:149:PRO:CD	2.77	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:207:VAL:HG22	2:B:221:LEU:HD23	1.96	0.48
2:B:120:LEU:HG	2:B:161:VAL:HG23	1.95	0.47
2:B:117:ASN:ND2	2:B:161:VAL:O	2.48	0.47
2:B:77:TYR:CD1	2:B:77:TYR:C	2.88	0.47
2:B:308:GLN:HA	2:B:308:GLN:OE1	2.14	0.47
2:B:74:ARG:CZ	2:B:90:PRO:HD3	2.45	0.46
2:B:36:ARG:NH1	2:B:36:ARG:CG	2.75	0.46
2:B:119:ILE:CG2	2:B:158:ILE:HG23	2.46	0.45
2:B:313:VAL:HG21	2:B:319:VAL:HG21	1.99	0.45
1:A:182:GLY:O	2:B:180:ASN:HB3	2.18	0.44
2:B:77:TYR:CD1	2:B:77:TYR:O	2.70	0.44
2:B:117:ASN:ND2	2:B:162:GLY:HA2	2.33	0.43
2:B:116:GLY:O	2:B:164:VAL:HG22	2.18	0.43
1:A:245:LYS:HE2	1:A:249:VAL:CG2	2.49	0.43
1:A:206:PRO:HB2	1:A:251:VAL:CG1	2.49	0.43
1:A:190:PHE:CZ	1:A:213:ILE:HG23	2.53	0.43
2:B:221:LEU:HD13	2:B:234:LEU:HD21	2.00	0.43
2:B:50:ILE:HG22	2:B:55:VAL:HG23	2.02	0.42
2:B:205:LEU:HG	2:B:274:CYS:HB2	2.01	0.42
2:B:6:PRO:HG2	2:B:88:SER:HB3	2.02	0.42
2:B:381:LYS:HB3	2:B:384:LEU:CD2	2.47	0.42
2:B:79:SER:HB3	2:B:82:ALA:HB3	2.02	0.42
1:A:197:PHE:HZ	1:A:216:GLU:HG2	1.84	0.42
2:B:118:VAL:HG12	2:B:161:VAL:HB	2.01	0.42
1:A:185:GLY:HA3	1:A:231:LEU:HG	2.02	0.41
1:A:197:PHE:CZ	1:A:216:GLU:HG2	2.54	0.41
2:B:331:THR:HG22	2:B:347:ARG:HA	2.01	0.41
2:B:223:CYS:O	2:B:256:GLN:HA	2.19	0.41
2:B:305:LEU:HD13	2:B:374:CYS:HB3	2.02	0.41
2:B:35:TYR:O	2:B:36:ARG:HG2	2.21	0.41
2:B:94:VAL:HG11	2:B:185:TRP:CD2	2.56	0.41
2:B:365:THR:H	2:B:368:HIS:HB2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/129 (76%)	94 (96%)	4 (4%)	0	100	100
2	B	394/437 (90%)	371 (94%)	23 (6%)	0	100	100
All	All	492/566 (87%)	465 (94%)	27 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/113 (79%)	86 (97%)	3 (3%)	37	72
2	B	337/370 (91%)	318 (94%)	19 (6%)	21	56
All	All	426/483 (88%)	404 (95%)	22 (5%)	23	59

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	209	LEU
1	A	214	LEU
1	A	233	ARG
2	B	3	LEU
2	B	5	LYS
2	B	42	LYS
2	B	76	TYR
2	B	77	TYR
2	B	141	HIS
2	B	148	GLN
2	B	191	LEU
2	B	205	LEU
2	B	219	LEU
2	B	240	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	245	LEU
2	B	267	SER
2	B	280	LEU
2	B	307	VAL
2	B	318	ASN
2	B	341	ASP
2	B	346	LEU
2	B	394	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	18	GLN
2	B	33	GLN
2	B	69	HIS
2	B	117	ASN
2	B	148	GLN
2	B	256	GLN
2	B	368	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	3,2	14,14,15	0.54	0	17,19,21	1.86	2 (11%)
3	NAG	C	2	3	14,14,15	0.38	0	17,19,21	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,2	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-O5-C5	6.33	120.77	112.19
3	C	1	NAG	C3-C4-C5	2.76	115.16	110.24

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	2	NAG	O5-C5-C6-O6
3	C	2	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	B	501	2	14,14,15	0.35	0	17,19,21	0.91	2 (11%)
4	NAG	B	502	2	14,14,15	0.36	0	17,19,21	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	501	2	-	0/6/23/26	0/1/1/1
4	NAG	B	502	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	NAG	C1-O5-C5	2.33	115.34	112.19
4	B	501	NAG	C3-C4-C5	2.03	113.85	110.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	100/129 (77%)	0.09	0	100 100	81, 121, 172, 249	0
2	B	396/437 (90%)	0.20	5 (1%)	77 51	80, 110, 174, 196	0
All	All	496/566 (87%)	0.18	5 (1%)	82 59	80, 112, 174, 249	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	33	GLN	5.6
2	B	80	ASP	4.3
2	B	308	GLN	3.2
2	B	30	GLN	2.3
2	B	152	ARG	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	1	14/15	0.45	0.29	171,180,183,184	0
3	NAG	C	2	14/15	0.85	0.45	184,186,186,186	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	B	502	14/15	0.77	0.18	179,181,184,184	0
4	NAG	B	501	14/15	0.82	0.16	153,156,161,162	0

6.5 Other polymers [i](#)

There are no such residues in this entry.