



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 3, 2024 – 01:00 PM JST

PDB ID : 4ZHT  
Title : Crystal structure of UDP-GlcNAc 2-epimerase  
Authors : Chen, S.C.; Yang, C.S.; Ko, T.P.; Chen, Y.  
Deposited on : 2015-04-27  
Resolution : 2.69 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	411	
1	B	411	
1	C	411	
1	D	411	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12681 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	384	Total	C	N	O	S	Se	0	0	0
			3026	1917	536	551	10	12			
1	B	382	Total	C	N	O	S	Se	0	0	0
			3011	1908	533	548	10	12			
1	C	383	Total	C	N	O	S	Se	0	0	0
			3016	1907	536	551	10	12			
1	D	383	Total	C	N	O	S	Se	0	0	0
			3018	1913	534	549	10	12			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	406	HIS	-	expression tag	UNP Q9Y223
A	407	HIS	-	expression tag	UNP Q9Y223
A	408	HIS	-	expression tag	UNP Q9Y223
A	409	HIS	-	expression tag	UNP Q9Y223
A	410	HIS	-	expression tag	UNP Q9Y223
A	411	HIS	-	expression tag	UNP Q9Y223
B	406	HIS	-	expression tag	UNP Q9Y223
B	407	HIS	-	expression tag	UNP Q9Y223
B	408	HIS	-	expression tag	UNP Q9Y223
B	409	HIS	-	expression tag	UNP Q9Y223
B	410	HIS	-	expression tag	UNP Q9Y223
B	411	HIS	-	expression tag	UNP Q9Y223
C	406	HIS	-	expression tag	UNP Q9Y223
C	407	HIS	-	expression tag	UNP Q9Y223
C	408	HIS	-	expression tag	UNP Q9Y223
C	409	HIS	-	expression tag	UNP Q9Y223
C	410	HIS	-	expression tag	UNP Q9Y223
C	411	HIS	-	expression tag	UNP Q9Y223
D	406	HIS	-	expression tag	UNP Q9Y223
D	407	HIS	-	expression tag	UNP Q9Y223

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	408	HIS	-	expression tag	UNP Q9Y223
D	409	HIS	-	expression tag	UNP Q9Y223
D	410	HIS	-	expression tag	UNP Q9Y223
D	411	HIS	-	expression tag	UNP Q9Y223

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula:  $C_9H_{14}N_2O_{12}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	C	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	D	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

- Molecule 3 is CYTIDINE-5'-MONOPHOSPHATE-5-N-ACETYLNEURAMINIC ACID (three-letter code: NCC) (formula:  $C_{20}H_{31}N_4O_{16}P$ ).



- 
- Chemical structure of BM7, a bicyclic molecule. The structure features a cyclohexane ring with several substituents and stereocenters. The ring carbons are labeled C1(R), C2(S), C3(R), C4(S), C5(R), and C6. Substituents include a hydroxyl group (O1, HO) at C1, a hydroxyl group (O3, OH) at C3, a hydroxyl group (O4, OH) at C4, a hydroxyl group (O5, OH) at C5, and a hydroxyl group (O6, OH) at C6. An amide group (N2, HN) is attached to C2, with a carbonyl group (C7, O7) and a methyl group (C8) also attached to C7. The stereochemistry is indicated by (R) and (S) labels and wedge/dash bonds.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	N	O	0	0
			15	8	1	6		


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	101	Total	O	0	0
			101	101		
5	B	78	Total	O	0	0
			78	78		
5	C	98	Total	O	0	0
			98	98		
5	D	54	Total	O	0	0
			54	54		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bifunctional UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase

Chain A: 





- Molecule 1: Bifunctional UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase

Chain B: 





- Molecule 1: Bifunctional UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase

Chain C: 



- Molecule 1: Bifunctional UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase

Chain D: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.04Å 98.10Å 154.72Å 90.00° 96.05° 90.00°	Depositor
Resolution (Å)	30.00 – 2.69 30.00 – 2.69	Depositor EDS
% Data completeness (in resolution range)	99.4 (30.00-2.69) 99.5 (30.00-2.69)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.51 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, $R_{free}$	0.189 , 0.225 0.194 , 0.227	Depositor DCC
$R_{free}$ test set	2183 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.8	Xtriage
Anisotropy	0.297	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 33.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12681	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BM7, UDP, NCC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	0/3071	0.87	10/4126 (0.2%)
1	B	0.54	0/3055	0.84	7/4103 (0.2%)
1	C	0.56	0/3059	0.83	7/4109 (0.2%)
1	D	0.52	0/3063	0.82	6/4115 (0.1%)
All	All	0.54	0/12248	0.84	30/16453 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	8	ARG	NE-CZ-NH1	8.73	124.67	120.30
1	D	177	ARG	NE-CZ-NH1	8.36	124.48	120.30
1	C	177	ARG	NE-CZ-NH1	8.20	124.40	120.30
1	A	177	ARG	NE-CZ-NH1	8.02	124.31	120.30
1	B	177	ARG	NE-CZ-NH1	8.00	124.30	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	192	ALA	Peptide
1	C	192	ALA	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3026	0	3066	24	0
1	B	3011	0	3053	24	0
1	C	3016	0	3056	19	0
1	D	3018	0	3060	20	0
2	A	25	0	11	0	0
2	B	25	0	11	0	0
2	C	25	0	11	0	0
2	D	25	0	11	0	0
3	A	41	0	29	0	0
3	B	41	0	29	0	0
3	C	41	0	29	0	0
3	D	41	0	29	0	0
4	C	15	0	0	0	0
5	A	101	0	0	4	0
5	B	78	0	0	5	0
5	C	98	0	0	3	0
5	D	54	0	0	2	0
All	All	12681	0	12395	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:ARG:NH2	1:A:105:ASP:OD2	2.07	0.87
1:C:59:ARG:NH2	5:C:601:HOH:O	2.08	0.86
1:A:380:GLN:NE2	5:A:602:HOH:O	2.10	0.85
1:B:60:MSE:HE2	1:B:281:HIS:CD2	2.18	0.79
1:D:354:GLN:O	5:D:601:HOH:O	2.01	0.78

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/411 (93%)	365 (96%)	12 (3%)	5 (1%)	10	26
1	B	380/411 (92%)	364 (96%)	12 (3%)	4 (1%)	12	30
1	C	381/411 (93%)	363 (95%)	15 (4%)	3 (1%)	16	38
1	D	381/411 (93%)	363 (95%)	13 (3%)	5 (1%)	10	26
All	All	1524/1644 (93%)	1455 (96%)	52 (3%)	17 (1%)	12	30

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	381	GLU
1	B	381	GLU
1	C	193	LYS
1	C	196	ASP
1	C	381	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	334/347 (96%)	324 (97%)	10 (3%)	36	65
1	B	332/347 (96%)	320 (96%)	12 (4%)	30	59
1	C	333/347 (96%)	324 (97%)	9 (3%)	40	69

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	333/347 (96%)	321 (96%)	12 (4%)	30	59
All	All	1332/1388 (96%)	1289 (97%)	43 (3%)	34	63

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	246	ARG
1	D	205	LEU
1	C	338	ASP
1	D	145	SER
1	D	246	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	127	ASN
1	B	127	ASN
1	C	127	ASN
1	C	157	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	UDP	B	501	-	24,26,26	1.17	3 (12%)	37,40,40	1.55	6 (16%)
3	NCC	A	502	-	40,43,43	0.78	0	49,65,65	1.43	7 (14%)
3	NCC	C	502	-	40,43,43	0.74	0	49,65,65	1.68	11 (22%)
2	UDP	C	501	-	24,26,26	1.20	3 (12%)	37,40,40	1.72	6 (16%)
4	BM7	C	503	-	15,15,15	1.15	1 (6%)	21,21,21	3.66	4 (19%)
3	NCC	D	502	-	40,43,43	0.79	0	49,65,65	1.38	5 (10%)
2	UDP	A	501	-	24,26,26	1.11	2 (8%)	37,40,40	1.69	5 (13%)
3	NCC	B	502	-	40,43,43	0.73	0	49,65,65	1.58	8 (16%)
2	UDP	D	501	-	24,26,26	1.12	3 (12%)	37,40,40	1.57	5 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	B	501	-	-	0/16/32/32	0/2/2/2
3	NCC	A	502	-	-	3/30/70/70	0/3/3/3
3	NCC	C	502	-	-	5/30/70/70	0/3/3/3
2	UDP	C	501	-	-	5/16/32/32	0/2/2/2
4	BM7	C	503	-	-	5/6/26/26	0/1/1/1
3	NCC	D	502	-	-	3/30/70/70	0/3/3/3
2	UDP	A	501	-	-	1/16/32/32	0/2/2/2
3	NCC	B	502	-	-	3/30/70/70	0/3/3/3
2	UDP	D	501	-	-	7/16/32/32	0/2/2/2

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	503	BM7	C1-C2	3.27	1.56	1.52
2	C	501	UDP	C2-N1	2.80	1.42	1.38
2	B	501	UDP	C4-N3	-2.52	1.34	1.38
2	A	501	UDP	C2-N1	2.38	1.42	1.38
2	A	501	UDP	C5-C4	-2.27	1.38	1.43

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	503	BM7	C1-C2-N2	11.91	124.53	110.73
4	C	503	BM7	O5-C1-C2	9.98	119.55	109.52
3	B	502	NCC	C18-C17-N5	4.79	124.21	116.10
2	C	501	UDP	C4-N3-C2	-4.73	120.35	126.58
3	C	502	NCC	C4-N3-C2	4.54	120.94	116.34

There are no chirality outliers.



5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	501	UDP	C5'-O5'-PA-O2A
2	C	501	UDP	C5'-O5'-PA-O3A
2	D	501	UDP	O4'-C4'-C5'-O5'
2	D	501	UDP	C5'-O5'-PA-O3A
3	A	502	NCC	C19-C16-N5-C17



There are no ring outliers.



No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	372/411 (90%)	-0.66	0 100 100	47, 66, 106, 163	0
1	B	370/411 (90%)	-0.53	0 100 100	43, 78, 117, 165	0
1	C	371/411 (90%)	-0.67	0 100 100	44, 63, 107, 150	0
1	D	371/411 (90%)	-0.42	0 100 100	50, 89, 135, 187	0
All	All	1484/1644 (90%)	-0.57	0 100 100	43, 72, 122, 187	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	BM7	C	503	15/15	0.62	0.16	74,87,108,113	0
3	NCC	D	502	41/41	0.93	0.07	52,66,77,85	0
3	NCC	B	502	41/41	0.95	0.06	58,64,75,78	0
2	UDP	D	501	25/25	0.95	0.07	49,64,74,78	0

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NCC	A	502	41/41	0.95	0.06	41,53,69,75	0
3	NCC	C	502	41/41	0.96	0.06	38,46,53,70	0
2	UDP	A	501	25/25	0.97	0.05	39,52,60,65	0
2	UDP	B	501	25/25	0.97	0.05	51,63,68,72	0
2	UDP	C	501	25/25	0.98	0.05	41,47,52,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around NCC B 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UDP D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around NCC A 502:**




$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around NCC C 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around UDP A 501:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around UDP B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around UDP C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.