



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 27, 2025 – 06:09 PM JST

PDB ID : 8ZHW
Title : Structure of Mokola lyssavirus glycoprotein in post-fusion state
Authors : Lu, G.W.; Yang, F.L.; Lin, S.; Yang, J.; Ye, F.
Deposited on : 2024-05-11
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1370 (3.20-3.20)
Clashscore	180529	1497 (3.20-3.20)
Ramachandran outliers	177936	1479 (3.20-3.20)
Sidechain outliers	177891	1478 (3.20-3.20)
RSRZ outliers	164620	1371 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	<div> <div>3%</div> <div> <div></div> <div>54%</div> <div>21%</div> <div>•</div> <div>22%</div> </div> </div>
1	B	436	<div> <div>4%</div> <div> <div></div> <div>48%</div> <div>15%</div> <div>•</div> <div>35%</div> </div> </div>
1	C	436	<div> <div>3%</div> <div> <div></div> <div>49%</div> <div>16%</div> <div>•</div> <div>34%</div> </div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 7217 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	0	0	0
			2663	1699	454	487	23			
1	B	285	Total	C	N	O	S	0	0	0
			2275	1456	387	413	19			
1	C	286	Total	C	N	O	S	0	0	0
			2279	1452	388	419	20			


There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	75	GLY	-	linker	UNP P0C572
A	76	GLY	-	linker	UNP P0C572
A	77	SER	-	linker	UNP P0C572
A	78	GLY	-	linker	UNP P0C572
A	79	GLY	-	linker	UNP P0C572
A	121	GLY	-	linker	UNP P0C572
A	122	GLY	-	linker	UNP P0C572
A	123	SER	-	linker	UNP P0C572
A	124	GLY	-	linker	UNP P0C572
A	125	GLY	-	linker	UNP P0C572
A	437	HIS	-	expression tag	UNP P0C572
A	438	HIS	-	expression tag	UNP P0C572
A	439	HIS	-	expression tag	UNP P0C572
A	440	HIS	-	expression tag	UNP P0C572
A	441	HIS	-	expression tag	UNP P0C572
A	442	HIS	-	expression tag	UNP P0C572
B	79	GLY	-	linker	UNP P0C572
B	80	GLY	-	linker	UNP P0C572
B	81	SER	-	linker	UNP P0C572
B	82	GLY	-	linker	UNP P0C572
B	83	GLY	-	linker	UNP P0C572
B	121	GLY	-	linker	UNP P0C572
B	122	GLY	-	linker	UNP P0C572

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	123	SER	-	linker	UNP P0C572
B	124	GLY	-	linker	UNP P0C572
B	125	GLY	-	linker	UNP P0C572
B	437	HIS	-	expression tag	UNP P0C572
B	438	HIS	-	expression tag	UNP P0C572
B	439	HIS	-	expression tag	UNP P0C572
B	440	HIS	-	expression tag	UNP P0C572
B	441	HIS	-	expression tag	UNP P0C572
B	442	HIS	-	expression tag	UNP P0C572
C	79	GLY	-	linker	UNP P0C572
C	80	GLY	-	linker	UNP P0C572
C	81	SER	-	linker	UNP P0C572
C	82	GLY	-	linker	UNP P0C572
C	83	GLY	-	linker	UNP P0C572
C	121	GLY	-	linker	UNP P0C572
C	122	GLY	-	linker	UNP P0C572
C	123	SER	-	linker	UNP P0C572
C	124	GLY	-	linker	UNP P0C572
C	125	GLY	-	linker	UNP P0C572
C	437	HIS	-	expression tag	UNP P0C572
C	438	HIS	-	expression tag	UNP P0C572
C	439	HIS	-	expression tag	UNP P0C572
C	440	HIS	-	expression tag	UNP P0C572
C	441	HIS	-	expression tag	UNP P0C572
C	442	HIS	-	expression tag	UNP P0C572



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.23Å 87.52Å 111.19Å 90.00° 111.87° 90.00°	Depositor
Resolution (Å)	48.37 – 3.20 48.37 – 3.20	Depositor EDS
% Data completeness (in resolution range)	95.0 (48.37-3.20) 95.0 (48.37-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 3.19Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.246 , 0.288 0.247 , 0.292	Depositor DCC
R_{free} test set	1493 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	107.2	Xtriage
Anisotropy	0.428	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 78.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7217	wwPDB-VP
Average B, all atoms (Å ²)	138.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/2720	0.52	2/3669 (0.1%)
1	B	0.29	0/2325	0.51	0/3132
1	C	0.28	0/2326	0.53	1/3133 (0.0%)
All	All	0.29	0/7371	0.52	3/9934 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	339	LEU	CA-CB-CG	7.82	133.28	115.30
1	A	252	CYS	N-CA-C	6.52	128.60	111.00
1	A	252	CYS	CB-CA-C	-5.35	99.70	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2663	0	2639	72	0
1	B	2275	0	2278	38	0
1	C	2279	0	2268	47	0
All	All	7217	0	7185	155	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 155 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:339:LEU:H	1:A:339:LEU:HD12	1.40	0.87
1:A:340:PRO:HG2	1:A:346:LYS:HZ1	1.39	0.86
1:A:340:PRO:HG2	1:A:346:LYS:NZ	1.91	0.85
1:C:339:LEU:HB2	1:C:372:LEU:HD22	1.63	0.80
1:B:1:GLU:HG3	1:B:331:ARG:HH12	1.47	0.79

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/436 (75%)	301 (92%)	27 (8%)	0	100	100
1	B	277/436 (64%)	256 (92%)	21 (8%)	0	100	100
1	C	276/436 (63%)	261 (95%)	15 (5%)	0	100	100
All	All	881/1308 (67%)	818 (93%)	63 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/388 (77%)	271 (91%)	28 (9%)	7	28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	257/388 (66%)	240 (93%)	17 (7%)	14	45
1	C	258/388 (66%)	245 (95%)	13 (5%)	20	54
All	All	814/1164 (70%)	756 (93%)	58 (7%)	12	42

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	13	LYS
1	C	364	ILE
1	B	265	LEU
1	C	344	CYS
1	C	215	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	27	ASN
1	C	27	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	338/436 (77%)	0.21	14 (4%)	42	28	85, 125, 204, 261	0
1	B	285/436 (65%)	0.27	19 (6%)	25	17	96, 136, 199, 232	0
1	C	286/436 (65%)	0.33	12 (4%)	41	28	81, 130, 201, 271	0
All	All	909/1308 (69%)	0.27	45 (4%)	35	24	81, 130, 203, 271	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	162	VAL	4.6
1	A	50	TYR	4.4
1	A	167	PRO	4.3
1	C	300	ARG	4.3
1	B	341	SER	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.