



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 15, 2024 – 06:35 AM EST

PDB ID : 1ZL0  
Title : Structure of Protein of Unknown Function PA5198 from Pseudomonas aeruginosa  
Authors : Osipiuk, J.; Evdokimova, E.; Savchenko, A.; Edwards, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2005-05-04  
Resolution : 1.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.21  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.004 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1365 (1.12-1.08)
Clashscore	180529	1561 (1.12-1.08)
Ramachandran outliers	177936	1524 (1.12-1.08)
Sidechain outliers	177891	1520 (1.12-1.08)
RSRZ outliers	164620	1365 (1.12-1.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>%</span> <span>82%</span> <span>14%</span> <span>..</span> </div> </div>
1	B	311	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <span>%</span> <span>83%</span> <span>13%</span> <span>..</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	TLA	A	1401	-	-	X	-
5	EDO	A	1408[B]	-	-	X	-
5	EDO	B	1411[A]	-	-	X	-
5	EDO	B	1411[B]	-	-	X	-
5	EDO	B	1412[B]	-	-	X	-
5	EDO	B	1413[A]	-	-	X	-
5	EDO	B	1413[B]	-	-	X	-
6	GOL	A	1402	-	X	X	-
7	PEG	B	1403[A]	-	-	X	-
7	PEG	B	1403[B]	-	-	X	-

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6196 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hypothetical protein PA5198.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	0	38	0
			2545	1599	474	463	9			
1	B	303	Total	C	N	O	S	0	40	0
			2503	1581	453	459	10			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	cloning artifact	UNP Q9HTZ1
A	0	HIS	-	cloning artifact	UNP Q9HTZ1
A	308	GLY	-	cloning artifact	UNP Q9HTZ1
A	309	SER	-	cloning artifact	UNP Q9HTZ1
B	-1	GLY	-	cloning artifact	UNP Q9HTZ1
B	0	HIS	-	cloning artifact	UNP Q9HTZ1
B	308	GLY	-	cloning artifact	UNP Q9HTZ1
B	309	SER	-	cloning artifact	UNP Q9HTZ1


- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	K	0	0
			1	1		
2	B	1	Total	K	0	0
			1	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	4	6		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	1
			8	4	4		
5	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	1
			8	4	4		
5	B	1	Total	C	O	0	1
			8	4	4		
5	B	1	Total	C	O	0	1
			8	4	4		
5	B	1	Total	C	O	0	1
			8	4	4		
5	B	1	Total	C	O	0	1
			8	4	4		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	1
			14	8	6		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	512	Total	O	0	2
			514	514		
8	B	523	Total	O	0	2
			525	525		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: hypothetical protein PA5198



- Molecule 1: hypothetical protein PA5198





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.68Å 79.80Å 70.83Å 90.00° 104.01° 90.00°	Depositor
Resolution (Å)	34.50 – 1.10 34.50 – 1.10	Depositor EDS
% Data completeness (in resolution range)	81.4 (34.50-1.10) 81.4 (34.50-1.10)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.16 (at 1.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.099 , 0.121 0.103 , 0.125	Depositor DCC
$R_{free}$ test set	4652 reflections (2.45%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	8.8	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6196	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, K, GOL, PEG, EDO, TLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.82	6/2737 (0.2%)	1.02	15/3713 (0.4%)
1	B	0.84	5/2702 (0.2%)	0.96	14/3668 (0.4%)
All	All	0.83	11/5439 (0.2%)	0.99	29/7381 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	225[A]	ARG	CZ-NH2	-8.59	1.21	1.33
1	A	225[B]	ARG	CZ-NH2	-8.59	1.21	1.33
1	B	88[A]	GLY	C-N	-8.52	1.14	1.34
1	B	88[B]	GLY	C-N	-8.52	1.14	1.34
1	A	64[A]	GLU	CD-OE2	7.42	1.33	1.25

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	225[A]	ARG	NE-CZ-NH2	-10.75	114.93	120.30
1	A	225[B]	ARG	NE-CZ-NH2	-10.75	114.93	120.30
1	B	15	ASP	CB-CG-OD2	-10.73	108.65	118.30
1	A	225[A]	ARG	NE-CZ-NH1	9.66	125.13	120.30
1	A	225[B]	ARG	NE-CZ-NH1	9.66	125.13	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	88[B]	GLY	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2545	0	2480	94	0
1	B	2503	0	2445	86	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
4	A	10	0	3	4	0
5	A	20	0	30	8	0
5	B	56	0	83	55	0
6	A	6	0	8	8	0
7	B	14	0	19	36	0
8	A	514	0	0	52	4
8	B	525	0	0	40	3
All	All	6196	0	5068	213	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 213 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:1413[B]:EDO:C2	7:B:1403[B]:PEG:H21	1.56	1.35
5:B:1413[B]:EDO:C1	7:B:1403[B]:PEG:C1	2.06	1.32
1:B:115[B]:SER:OG	5:B:1412[B]:EDO:H12	1.28	1.30
1:A:228[A]:ARG:NH1	8:A:1898:HOH:O	1.61	1.28
1:A:50[B]:ARG:NH1	8:A:1844:HOH:O	1.62	1.26

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:1815:HOH:O	8:B:1438:HOH:O[1_554]	1.66	0.54
8:A:1797:HOH:O	8:A:1800:HOH:O[1_455]	2.09	0.11
8:B:1839:HOH:O	8:B:1844:HOH:O[2_657]	2.09	0.11
8:A:1685:HOH:O	8:A:1899:HOH:O[1_455]	2.12	0.08
8:A:1808:HOH:O	8:B:1681:HOH:O[1_455]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	341/311 (110%)	333 (98%)	7 (2%)	1 (0%)	37	13
1	B	340/311 (109%)	334 (98%)	6 (2%)	0	100	100
All	All	681/622 (110%)	667 (98%)	13 (2%)	1 (0%)	44	15

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/242 (114%)	272 (99%)	3 (1%)	70	35
1	B	274/242 (113%)	271 (99%)	3 (1%)	70	35
All	All	549/484 (113%)	543 (99%)	6 (1%)	70	35

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	257[A]	GLU
1	B	257[B]	GLU
1	B	279	HIS
1	A	58	TYR
1	A	4	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	175	GLN
1	A	279	HIS
1	B	71	HIS
1	A	71	HIS
1	A	47	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 3 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EDO	B	1414[B]	-	3,3,3	0.46	0	2,2,2	0.54	0
5	EDO	B	1413[A]	-	3,3,3	0.63	0	2,2,2	0.88	0
5	EDO	B	1406[B]	-	3,3,3	0.25	0	2,2,2	0.61	0
6	GOL	A	1402	-	5,5,5	0.96	0	5,5,5	3.33	4 (80%)
5	EDO	B	1411[B]	-	3,3,3	0.31	0	2,2,2	1.32	0
5	EDO	B	1412[A]	-	3,3,3	0.48	0	2,2,2	0.40	0
5	EDO	B	1404	-	3,3,3	0.29	0	2,2,2	0.32	0
5	EDO	B	1413[B]	7	3,3,3	0.63	0	2,2,2	1.22	0
5	EDO	A	1407	-	3,3,3	0.42	0	2,2,2	0.63	0
5	EDO	B	1412[B]	-	3,3,3	0.37	0	2,2,2	1.88	1 (50%)
5	EDO	A	1410	-	3,3,3	0.32	0	2,2,2	0.30	0
5	EDO	A	1408[A]	-	3,3,3	1.13	0	2,2,2	1.16	0
5	EDO	A	1415	-	3,3,3	0.53	0	2,2,2	0.31	0
5	EDO	B	1405	-	3,3,3	0.85	0	2,2,2	1.01	0
7	PEG	B	1403[A]	-	6,6,6	0.42	0	5,5,5	2.92	3 (60%)
5	EDO	B	1409[A]	-	3,3,3	0.60	0	2,2,2	0.12	0
5	EDO	A	1408[B]	-	3,3,3	0.86	0	2,2,2	1.01	0
5	EDO	B	1414[A]	-	3,3,3	0.31	0	2,2,2	1.14	0
7	PEG	B	1403[B]	5	6,6,6	0.89	0	5,5,5	1.01	0
5	EDO	B	1409[B]	-	3,3,3	0.71	0	2,2,2	0.53	0
4	TLA	A	1401	2	9,9,9	1.59	1 (11%)	12,12,12	1.48	3 (25%)
5	EDO	B	1406[A]	-	3,3,3	0.38	0	2,2,2	0.91	0
5	EDO	B	1411[A]	-	3,3,3	0.37	0	2,2,2	2.76	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	1414[B]	-	-	0/1/1/1	-
5	EDO	B	1413[A]	-	-	1/1/1/1	-
5	EDO	B	1406[B]	-	-	1/1/1/1	-
6	GOL	A	1402	-	-	4/4/4/4	-
5	EDO	B	1411[B]	-	-	1/1/1/1	-
5	EDO	B	1412[A]	-	-	1/1/1/1	-
5	EDO	B	1404	-	-	0/1/1/1	-
5	EDO	B	1413[B]	7	-	0/1/1/1	-
5	EDO	A	1407	-	-	0/1/1/1	-
5	EDO	B	1412[B]	-	-	1/1/1/1	-
5	EDO	A	1410	-	-	0/1/1/1	-

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	1408[A]	-	-	1/1/1/1	-
5	EDO	A	1415	-	-	1/1/1/1	-
5	EDO	B	1405	-	-	0/1/1/1	-
7	PEG	B	1403[A]	-	-	2/4/4/4	-
5	EDO	B	1409[A]	-	-	0/1/1/1	-
5	EDO	A	1408[B]	-	-	0/1/1/1	-
5	EDO	B	1414[A]	-	-	1/1/1/1	-
7	PEG	B	1403[B]	5	-	3/4/4/4	-
5	EDO	B	1409[B]	-	-	1/1/1/1	-
4	TLA	A	1401	2	-	0/12/12/12	-
5	EDO	B	1406[A]	-	-	0/1/1/1	-
5	EDO	B	1411[A]	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1401	TLA	C2-C1	2.13	1.55	1.52

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	1403[A]	PEG	O2-C2-C1	4.68	130.74	110.11
6	A	1402	GOL	O3-C3-C2	-4.42	90.49	110.38
6	A	1402	GOL	O2-C2-C3	4.10	126.16	109.18
7	B	1403[A]	PEG	O1-C1-C2	-3.86	89.11	111.82
5	B	1411[A]	EDO	O1-C1-C2	-3.83	83.22	112.39

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1402	GOL	O1-C1-C2-C3
6	A	1402	GOL	C1-C2-C3-O3
7	B	1403[B]	PEG	O2-C3-C4-O4
6	A	1402	GOL	O2-C2-C3-O3
5	B	1406[B]	EDO	O1-C1-C2-O2

There are no ring outliers.

17 monomers are involved in 84 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1413[A]	EDO	6	0
5	B	1406[B]	EDO	1	0
6	A	1402	GOL	8	0
5	B	1411[B]	EDO	5	0
5	B	1412[A]	EDO	2	0
5	B	1413[B]	EDO	17	0
5	A	1407	EDO	3	0
5	B	1412[B]	EDO	18	0
5	A	1410	EDO	1	0
5	A	1408[A]	EDO	2	0
5	A	1415	EDO	1	0
7	B	1403[A]	PEG	10	0
5	A	1408[B]	EDO	4	0
7	B	1403[B]	PEG	26	0
5	B	1409[B]	EDO	2	0
4	A	1401	TLA	4	0
5	B	1411[A]	EDO	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	88[A]:GLY	C	89:TYR	N	1.14



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	306/311 (98%)	-0.26	4 (1%) 74 79	3, 9, 18, 31	39 (12%)
1	B	303/311 (97%)	-0.49	2 (0%) 84 86	4, 7, 13, 23	41 (13%)
All	All	609/622 (97%)	-0.37	6 (0%) 79 82	3, 8, 16, 31	80 (13%)

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	308[A]	GLY	3.2
1	A	4	ARG	3.2
1	A	3	SER	3.1
1	B	7[A]	SER	2.5
1	A	254	PRO	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	EDO	B	1413[A]	4/4	0.77	0.13	11,15,17,20	4
5	EDO	B	1413[B]	4/4	0.77	0.13	12,12,15,18	4
7	PEG	B	1403[A]	7/7	0.84	0.12	8,13,18,23	7
7	PEG	B	1403[B]	7/7	0.84	0.12	8,16,19,21	7
5	EDO	A	1408[A]	4/4	0.86	0.12	10,16,18,26	4
5	EDO	A	1408[B]	4/4	0.86	0.12	6,14,14,20	4
5	EDO	B	1414[A]	4/4	0.90	0.11	13,17,18,19	4
5	EDO	B	1414[B]	4/4	0.90	0.11	10,18,19,23	4
5	EDO	B	1409[A]	4/4	0.90	0.09	17,19,20,22	4
5	EDO	B	1409[B]	4/4	0.90	0.09	15,17,17,18	4
5	EDO	B	1406[A]	4/4	0.93	0.08	14,16,17,18	4
5	EDO	B	1406[B]	4/4	0.93	0.08	13,14,15,21	4
5	EDO	B	1411[A]	4/4	0.93	0.09	9,10,11,23	4
5	EDO	B	1411[B]	4/4	0.93	0.09	11,11,14,19	4
5	EDO	B	1412[A]	4/4	0.93	0.09	9,12,15,20	4
5	EDO	B	1412[B]	4/4	0.93	0.09	9,12,17,17	4
6	GOL	A	1402	6/6	0.94	0.10	11,12,13,19	6
5	EDO	A	1415	4/4	0.94	0.07	20,21,23,25	0
5	EDO	B	1404	4/4	0.94	0.08	13,15,17,17	0
5	EDO	A	1410	4/4	0.95	0.07	14,15,15,21	0
5	EDO	A	1407	4/4	0.96	0.07	11,12,17,18	0
4	TLA	A	1401	10/10	0.96	0.06	9,11,12,17	10
5	EDO	B	1405	4/4	0.96	0.07	13,17,18,21	0
3	NA	A	1418	1/1	0.99	0.03	10,10,10,10	0
2	K	A	1417	1/1	1.00	0.04	9,9,9,9	1
2	K	B	1416	1/1	1.00	0.02	8,8,8,8	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.