



# Full wwPDB NMR Structure Validation Report ⓘ

Mar 19, 2025 – 04:09 AM EDT

PDB ID : 1ZNT  
Title : 18 NMR structures of AcAMP2-Like Peptide with non Natural Fluoroaromatic Residue (AcAMP2F18Pff/Y20Pff) complex with N,N,N-triacetylchitotriose  
Authors : Chavez, M.I.; Andreu, C.; Vidal, P.; Aboitiz, N.; Freire, F.; Groves, P.; Asensio, J.L.; Asensio, G.; Muraki, M.; Canada, F.J.; Jimenez-Barbero, J.  
Deposited on : 2005-05-12

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	31	
2	B	3	

## 2 Ensemble composition and analysis

This entry contains 18 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:17, A:19-A:19, A:21-A:28 (24)	0.80	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 3, 4, 7, 9, 10, 11, 12, 14, 15, 16, 17
2	1, 13, 18
3	5, 8
Single-model clusters	6


### 3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 506 atoms, of which 245 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2.

Mol	Chain	Residues	Atoms							Trace
1	A	31	Total	C	F	H	N	O	S	1
			422	130	2	204	43	36	7	

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					Trace
2	B	3	Total	C	H	N	O	0
			84	24	41	3	16	

## 4 Residue-property plots [i](#)


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



NAG1  
NAG2  
NAG3

#### 4.2.2 Score per residue for model 2

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  61% 10% 23%

V1 G2 R6 G7 R8 C9 F10 S11 Q17 F18 G19 F20 G29 R30 NH231

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  100%

NAG1  
NAG2  
NAG3

#### 4.2.3 Score per residue for model 3

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  61% 13% 23%

V1 G2 E3 R6 G7 R8 C9 F18 G19 F20 P25 G29 R30 NH231

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  100%

NAG1  
NAG2  
NAG3

#### 4.2.4 Score per residue for model 4

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  58% 19% 23%

V1 G2 R6 C9 P10 S11 F18 G19 F20 P25 R26 Y27 C28 G29 R30 NH231

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  67% 33%

MAG1  
MAG2  
MAG3

#### 4.2.5 Score per residue for model 5

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  61% 10% 6% 23%

V1 G2 E3 R6 G7 R8 C9 F18 G19 F20 C21 G29 R30 NH231

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  100%

MAG1  
MAG2  
MAG3

#### 4.2.6 Score per residue for model 6

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  61% 13% 1% 23%

V1 G2 E3 C4 V5 R6 C9 F18 G19 F20 K28 G29 R30 NH231

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose


Chain B:  100%

MAG1  
MAG2  
MAG3

#### 4.2.7 Score per residue for model 7 (medoid)

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  68% 10% 23%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  100%



#### 4.2.8 Score per residue for model 8

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  58% 13% 23%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose


Chain B:  67% 33%



#### 4.2.9 Score per residue for model 9

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  58% 16% 23%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  67% 33%






#### 4.2.10 Score per residue for model 10

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A: 



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose


Chain B: 




#### 4.2.11 Score per residue for model 11

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A: 



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose


Chain B: 



#### 4.2.12 Score per residue for model 12

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A: 



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose


Chain B: 



#### 4.2.13 Score per residue for model 13

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A: 71% 6% 23%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose


Chain B: 100%



#### 4.2.14 Score per residue for model 14

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A: 61% 10% 6% 23%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose


Chain B: 100%



#### 4.2.15 Score per residue for model 15

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A: 58% 16% 23%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  100%

MAG1  
MAG2  
MAG3

#### 4.2.16 Score per residue for model 16

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  61% 13% 23%

V1 G2 R6 G7 R8 C9 F10 Q17 F18 G19 F20 Y27 C28 G29 R30 NH231

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  67% 33%

MAG1  
MAG2  
MAG3

#### 4.2.17 Score per residue for model 17

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  61% 16% 23%

V1 G2 V5 R6 G7 R8 S11 F18 G19 F20 K23 G29 R30 NH231

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose


Chain B:  100%

MAG1  
MAG2  
MAG3

#### 4.2.18 Score per residue for model 18

- Molecule 1: AMARANTHUS CAUDATUS ANTIMICROBIAL PEPTIDE 2

Chain A:  65% 13% 23%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  100%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *The structures are based on a total 314 cross peaks, 248 NOE-derived distance restraints, and finally 208 distance constraints and 18 come from cys-cys disulfide were used in the final round of calculation.*

Of the 30 calculated structures, 18 were deposited, based on the following criterion: *Fewest restraint violation, secondary lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	1.5
Amber	refinement	5.0

No chemical shift data was provided.

## 6 Model quality

### 6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PFF, NH2, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.57±0.01	0±0/168 ( 0.0± 0.0%)	1.13±0.10	1±1/219 ( 0.4± 0.3%)
All	All	0.57	0/3024 ( 0.0%)	1.13	16/3942 ( 0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.6±0.6
All	All	0	10

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	6	ARG	NE-CZ-NH1	10.42	125.51	120.30	8	4
1	A	8	ARG	NE-CZ-NH1	7.95	124.27	120.30	10	10
1	A	6	ARG	NE-CZ-NH2	5.52	123.06	120.30	4	2

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	6	ARG	Sidechain	4
1	A	8	ARG	Sidechain	3
1	A	27	TYR	Sidechain	3

## 6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	167	156	156	0±1
2	B	43	41	39	0±1
All	All	3780	3546	3510	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:27:TYR:CE1	2:B:3:NAG:H83	0.69	2.22	16	2
1:A:27:TYR:CZ	2:B:3:NAG:H83	0.43	2.48	8	3
1:A:27:TYR:CE2	2:B:3:NAG:H83	0.43	2.49	8	1
1:A:17:GLN:CD	1:A:17:GLN:H	0.40	2.19	2	1

## 6.3 Torsion angles

### 6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	24/31 (77%)	19±2 (77±7%)	4±2 (16±7%)	2±1 (7±5%)	<b>2</b>	<b>15</b>
All	All	432/558 (77%)	334 (77%)	67 (16%)	31 (7%)	<b>2</b>	<b>15</b>

All 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	11	SER	8
1	A	10	PRO	7
1	A	5	VAL	6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	25	PRO	3
1	A	3	GLU	2
1	A	4	CYS	2
1	A	6	ARG	1
1	A	7	GLY	1
1	A	9	CYS	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	19/21 (90%)	16±1 (86±7%)	3±1 (14±7%)	5	44
All	All	342/378 (90%)	293 (86%)	49 (14%)	5	44

All 9 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	9	CYS	13
1	A	6	ARG	12
1	A	8	ARG	7
1	A	3	GLU	5
1	A	26	LYS	3
1	A	23	LYS	3
1	A	21	CYS	2
1	A	28	CYS	2
1	A	17	GLN	2

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.



In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	PFF	A	18	1	11,12,13	1.18±0.02	0±0 (1±3%)
1	PFF	A	20	1	11,12,13	1.14±0.04	0±0 (0±2%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	PFF	A	18	1	10,15,17	1.21±0.04	1±0 (10±2%)
1	PFF	A	20	1	10,15,17	1.16±0.04	1±0 (10±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PFF	A	18	1	-	0±0,5,6,8	0±0,1,1,1
1	PFF	A	20	1	-	0±0,5,6,8	0±0,1,1,1

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	18	PFF	CE2-CZ	2.03	1.41	1.37	12	2
1	A	18	PFF	CE1-CZ	2.01	1.41	1.37	15	1
1	A	20	PFF	CE1-CZ	2.01	1.41	1.37	4	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	20	PFF	CE2-CZ-CE1	2.35	119.72	122.80	18	18
1	A	18	PFF	CE2-CZ-CE1	2.34	119.73	122.80	11	18
1	A	18	PFF	CB-CG-CD2	2.10	116.99	120.90	15	1

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 6.5 Carbohydrates

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	NAG	B	1	2	15,15,15	0.77±0.07	0±0 (2±3%)
2	NAG	B	2	2	14,14,15	0.73±0.09	0±0 (0±0%)
2	NAG	B	3	2	14,14,15	0.54±0.08	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	NAG	B	1	2	21,21,21	1.35±0.25	3±1 (15±6%)
2	NAG	B	2	2	17,19,21	1.54±0.33	2±1 (14±6%)
2	NAG	B	3	2	17,19,21	1.33±0.14	2±1 (13±5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2	-	0±0,6,26,26	0±0,1,1,1
2	NAG	B	2	2	-	0±0,6,23,26	0±0,1,1,1
2	NAG	B	3	2	-	0±0,6,23,26	0±0,1,1,1

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	B	1	NAG	C2-N2	2.10	1.49	1.45	3	6

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	2	NAG	C2-N2-C7	5.81	130.69	122.90	6	12
2	B	1	NAG	C2-N2-C7	3.95	132.36	123.11	3	8
2	B	2	NAG	O5-C1-C2	3.87	105.31	111.29	15	8
2	B	1	NAG	C4-C3-C2	3.81	115.95	110.40	16	5
2	B	2	NAG	C1-C2-N2	3.69	116.25	110.43	8	8
2	B	2	NAG	C1-O5-C5	3.48	116.85	112.19	8	3
2	B	1	NAG	C1-C2-N2	3.46	106.72	110.73	5	5
2	B	3	NAG	O5-C5-C6	3.43	114.34	107.66	6	1
2	B	2	NAG	C3-C4-C5	3.33	104.19	110.23	6	7
2	B	1	NAG	O5-C1-C2	3.27	112.80	109.52	4	3
2	B	1	NAG	C3-C2-N2	3.23	116.58	110.62	8	7
2	B	1	NAG	C3-C4-C5	3.08	104.65	110.23	15	3
2	B	3	NAG	O3-C3-C4	2.92	103.49	110.38	12	13
2	B	3	NAG	O4-C4-C5	2.79	116.19	109.32	17	15
2	B	2	NAG	C4-C3-C2	2.74	115.03	111.02	5	5
2	B	1	NAG	O4-C4-C3	2.69	116.72	110.38	6	4
2	B	1	NAG	O3-C3-C2	2.67	104.27	109.58	6	3
2	B	1	NAG	O5-C5-C4	2.62	114.42	109.70	6	2
2	B	3	NAG	O5-C5-C4	2.48	104.80	110.83	7	2
2	B	3	NAG	C3-C4-C5	2.43	105.82	110.23	7	7
2	B	1	NAG	O5-C5-C6	2.38	112.35	106.44	17	2
2	B	1	NAG	C1-C2-C3	2.35	107.34	110.54	8	3
2	B	1	NAG	O1-C1-O5	2.29	103.60	110.41	7	11
2	B	2	NAG	O7-C7-N2	2.19	125.85	121.98	6	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	1	NAG	O7-C7-N2	2.18	125.83	121.98	8	1
2	B	3	NAG	C2-N2-C7	2.06	125.66	122.90	8	1
2	B	3	NAG	O7-C7-N2	2.03	125.57	121.98	8	1
2	B	3	NAG	O7-C7-C8	2.02	118.47	122.05	16	1


There are no chirality outliers.

All unique torsion outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
2	B	2	NAG	C1-C2-N2-C7	1

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided