



wwPDB NMR Structure Validation Summary Report ⓘ

May 28, 2025 – 12:11 PM JST

PDB ID : 8ZNQ / pdb_00008znq
BMRB ID : 36669
Title : Solution structure of the complex of naphthyridine-azaquinolone and an RNA with ACG/AUA motif
Authors : Fujiwara, A.; Chen, Q.; Nakatani, K.; Murata, A.; Kawai, G.
Deposited on : 2024-05-27

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.5 (274361), CSD as541be (2020)
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment is 24%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	210492	14027
RNA backbone	6643	756

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	30	 <div>47% 37% 17%</div>

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.


3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1019 atoms, of which 350 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (30-MER).

Mol	Chain	Residues	Atoms						Trace
1	A	30	Total	C	H	N	O	P	0
			959	284	324	110	212	29	

- Molecule 2 is N³-{3-[(7-METHYL-1,8-NAPHTHYRIDIN-2-YL)AMINO]-3-OXOPROPYL}-N¹ -[(7-OXO-7,8-DIHYDRO-1,8-NAPHTHYRIDIN-2-YL)METHYL]-BETA-ALANINAMIDE (CCD ID: NAZ) (formula: C₂₄H₂₅N₇O₃) (labeled as "Ligand of Interest" by depositor).




4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: RNA (30-MER)



4.2 Residue scores for the representative (author defined) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: RNA (30-MER)



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure calculation	1.3
CNS	refinement	1.3

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	168
Number of shifts mapped to atoms	144
Number of unparsed shifts	0
Number of shifts with mapping errors	24
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	24%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAZ

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	635	324	324	34±1
2	A	34	26	25	26±2
All	All	6690	3500	3490	348

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

5 of 62 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:7:A:N7	2:A:101:NAZ:H43	0.66	2.06	2	10
1:A:7:A:H62	2:A:101:NAZ:H38	0.64	1.52	3	5
1:A:7:A:C2	2:A:101:NAZ:C28	0.63	2.82	5	10
1:A:21:U:C5	2:A:101:NAZ:H42	0.63	2.28	5	9
2:A:101:NAZ:O20	2:A:101:NAZ:H45	0.61	1.94	9	5

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	29/30 (97%)	9±0 (31±0%)	0±0 (0±0%)	0.46±0.00
All	All	290/300 (97%)	90 (31%)	0 (0%)	0.46

The overall RNA backbone suiteness is 0.43.

5 of 9 unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	7	A	10
1	A	8	C	10
1	A	10	G	10
1	A	15	U	10
1	A	16	U	10

There are no RNA pucker outliers to report.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	NAZ	A	101	-	36,37,37	2.15±0.01	10±1 (26±1%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	NAZ	A	101	-	47,50,50	2.08±0.07	16±1 (32±1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAZ	A	101	-	-	0±0,17,17,17	0±0,4,4,4

5 of 11 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	101	NAZ	C22-N23	6.17	1.47	1.33	6	10
2	A	101	NAZ	O34-C24	5.82	1.39	1.24	6	10
2	A	101	NAZ	C7-N18	4.34	1.45	1.35	4	10
2	A	101	NAZ	C28-N26	4.01	1.43	1.35	7	10
2	A	101	NAZ	C31-C30	3.09	1.43	1.36	6	10

5 of 19 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.


Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	A	101	NAZ	C27-C29-C28	4.77	121.40	117.75	5	10
2	A	101	NAZ	C8-N18-C7	4.73	119.78	128.25	6	10
2	A	101	NAZ	C15-C13-C12	4.01	120.82	117.75	10	10
2	A	101	NAZ	C25-C27-C29	3.85	114.86	120.82	10	10
2	A	101	NAZ	C11-C13-C12	3.84	120.69	117.75	4	10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 24% for the well-defined parts and 24% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chemical_shifts_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	168
Number of shifts mapped to atoms	144
Number of unparsed shifts	0
Number of shifts with mapping errors	24
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. First 5 (of 24) occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	31	NAZ	H35	14.81	0.0009	1
1	A	31	NAZ	H36	4.433	0.0019	1
1	A	31	NAZ	H37	3.875	0.0017	1
1	A	31	NAZ	H38	3.677	0.000	1
1	A	31	NAZ	H39	3.503	0.000	1
1	A	31	NAZ	H41	3.391	0.0015	1
1	A	31	NAZ	H42	3.229	0.000	1
1	A	31	NAZ	H43	2.674	0.000	1
1	A	31	NAZ	H44	2.614	0.000	1
1	A	31	NAZ	H45	8.142	0.0019	1
1	A	31	NAZ	H46	7.806	0.0019	1
1	A	31	NAZ	H47	6.461	0.0036	1
1	A	31	NAZ	H48	7.183	0.0017	1
1	A	31	NAZ	H49	10.8	0.0029	1

Continued on next page...

Continued from previous page...

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	31	NAZ	H50	2.28	0.0007	1
1	A	31	NAZ	H51	2.28	0.0007	1
1	A	31	NAZ	H52	2.28	0.0007	1
1	A	31	NAZ	H53	2.951	0.000	1
1	A	31	NAZ	H54	2.88	0.000	1
1	A	31	NAZ	H55	8.687	0.0022	1
1	A	31	NAZ	H56	6.349	0.0016	1
1	A	31	NAZ	H57	7.389	0.0021	1
1	A	31	NAZ	H58	6.236	0.0016	1
1	A	31	NAZ	H59	6.828	0.0024	1

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 24%, i.e. 136 atoms were assigned a chemical shift out of a possible 564. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Sugar	76/330 (23%)	76/180 (42%)	0/150 (0%)	0/0 (—%)
Base	60/234 (26%)	60/144 (42%)	0/50 (0%)	0/40 (0%)
Overall	136/564 (24%)	136/324 (42%)	0/200 (0%)	0/40 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index*(RCI) plot could be generated from the current chemical shift list. RCI is only applicable to proteins

8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	440
Intra-residue ($ i-j =0$)	168
Sequential ($ i-j =1$)	100
Medium range ($ i-j >1$ and $ i-j <5$)	10
Long range ($ i-j \geq 5$)	74
Inter-chain	56
Hydrogen bond restraints	32
Disulfide bond restraints	0
Total dihedral-angle restraints	245
Number of unmapped restraints	95
Number of restraints per residue	22.1
Number of long range restraints per residue ¹	3.3

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	18.8	0.19
0.2-0.5 (Medium)	4.0	0.3
>0.5 (Large)	None	None

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	Max (°)
1.0-10.0 (Small)	1.2	1.83
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None

9 Distance violation analysis ⓘ


9.1 Summary of distance violations ⓘ

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	168	38.2	18	10.7	4.1	17	10.1	3.9
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	64	14.5	3	4.7	0.7	2	3.1	0.5
Sidechain-Sidechain	104	23.6	15	14.4	3.4	15	14.4	3.4
Sequential (i-j =1)	100	22.7	1	1.0	0.2	1	1.0	0.2
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	36	8.2	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	64	14.5	1	1.6	0.2	1	1.6	0.2
Medium range (i-j >1 & i-j <5)	10	2.3	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	4	0.9	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	6	1.4	0	0.0	0.0	0	0.0	0.0
Long range (i-j ≥5)	74	16.8	4	5.4	0.9	4	5.4	0.9
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	8	1.8	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	66	15.0	4	6.1	0.9	4	6.1	0.9
Inter-chain	56	12.7	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	9	2.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	47	10.7	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	32	7.3	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	440	100.0	23	5.2	5.2	22	5.0	5.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	121	27.5	3	2.5	0.7	2	1.7	0.5
Sidechain-Sidechain	319	72.5	20	6.3	4.5	20	6.3	4.5

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfid bonds are counted in their appropriate category on the x-axis


9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	18	1	0	4	0	23	0.17	0.3	0.06	0.16
2	18	1	0	4	0	23	0.17	0.3	0.06	0.16
3	18	1	0	4	0	23	0.17	0.3	0.06	0.16
4	18	1	0	4	0	23	0.17	0.3	0.05	0.16
5	17	1	0	4	0	22	0.17	0.3	0.06	0.16
6	18	1	0	4	0	23	0.17	0.3	0.06	0.16
7	18	1	0	4	0	23	0.17	0.3	0.06	0.16
8	18	1	0	4	0	23	0.17	0.3	0.05	0.16
9	17	1	0	4	0	22	0.17	0.3	0.06	0.16
10	18	1	0	4	0	23	0.17	0.3	0.06	0.16

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 385(IR:150, SQ:99, MR:10, LR:70, IC:56) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	0	0	0	0	0	1	10.0
0	0	0	0	0	0	2	20.0
0	0	0	0	0	0	3	30.0

Continued on next page...


Continued from previous page...

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	0	0	0	0	0	4	40.0
0	0	0	0	0	0	5	50.0
0	0	0	0	0	0	6	60.0
0	0	0	0	0	0	7	70.0
1	0	0	0	0	1	8	80.0
0	0	0	0	0	0	9	90.0
17	1	0	4	0	22	10	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations

9.3.1 Bar graph : Distance violation statistics for the ensemble ⓘ




9.4 Most violated distance restraints in the ensemble ⓘ

9.4.1 Histogram : Distribution of mean distance violations ⓘ

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models

in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,120)	1:3:A:C:H41	1:3:A:C:H6	10	0.3	0.0	0.3
(1,267)	1:24:A:A:H62	1:24:A:A:H61	10	0.3	0.0	0.3
(1,234)	1:20:A:C:H41	1:20:A:C:H6	10	0.28	0.0	0.28
(1,199)	1:12:A:C:H5	1:19:A:G:H1	10	0.22	0.0	0.22
(1,372)	1:15:A:U:H1'	1:15:A:U:H6	10	0.19	0.0	0.19
(1,388)	1:25:A:C:H41	1:25:A:C:H42	10	0.19	0.0	0.19
(1,210)	1:13:A:C:H5	1:18:A:G:H1	10	0.18	0.0	0.18
(1,237)	1:20:A:C:H42	1:20:A:C:H6	10	0.18	0.0	0.18
(1,380)	1:20:A:C:H2'	1:20:A:C:H6	10	0.18	0.0	0.18
(1,362)	1:9:A:G:H22	1:9:A:G:H1	10	0.17	0.0	0.17

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,267)	1:24:A:A:H62	1:24:A:A:H61	1	0.3
(1,267)	1:24:A:A:H62	1:24:A:A:H61	2	0.3
(1,267)	1:24:A:A:H62	1:24:A:A:H61	3	0.3
(1,267)	1:24:A:A:H62	1:24:A:A:H61	4	0.3
(1,267)	1:24:A:A:H62	1:24:A:A:H61	6	0.3
(1,267)	1:24:A:A:H62	1:24:A:A:H61	7	0.3
(1,267)	1:24:A:A:H62	1:24:A:A:H61	8	0.3
(1,267)	1:24:A:A:H62	1:24:A:A:H61	9	0.3
(1,267)	1:24:A:A:H62	1:24:A:A:H61	10	0.3
(1,120)	1:3:A:C:H41	1:3:A:C:H6	1	0.3

10 Dihedral-angle violation analysis ⓘ


10.1 Summary of dihedral-angle violations ⓘ

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
NU0	27	11.0	2	7.4	0.8	0	0.0	0.0
ALPHA	16	6.5	0	0.0	0.0	0	0.0	0.0
GAMMA	16	6.5	0	0.0	0.0	0	0.0	0.0
NU2	27	11.0	0	0.0	0.0	0	0.0	0.0
CHI	30	12.2	1	3.3	0.4	1	3.3	0.4
ZETA	16	6.5	0	0.0	0.0	0	0.0	0.0
EPSILON	16	6.5	0	0.0	0.0	0	0.0	0.0
NU1	27	11.0	0	0.0	0.0	0	0.0	0.0
NU3	27	11.0	0	0.0	0.0	0	0.0	0.0
BETA	16	6.5	0	0.0	0.0	0	0.0	0.0
NU4	27	11.0	0	0.0	0.0	0	0.0	0.0
Total	245	100.0	3	1.2	1.2	1	0.4	0.4

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations ⓘ




Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model ⓘ

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Number of violations												Mean (°)	Max (°)	SD (°)	%
	NU0	ALPHA	GAMMA	NU2	CHI	ZETA	EPSILON	NU1	NU3	BETA	NU4	Total				
1	0	0	0	0	1	0	0	0	0	0	0	1	1.44	1.44	0.0	3.7
2	0	0	0	0	1	0	0	0	0	0	0	1	1.72	1.72	0.0	3.7
3	0	0	0	0	1	0	0	0	0	0	0	1	1.52	1.52	0.0	3.7
4	0	0	0	0	1	0	0	0	0	0	0	1	1.64	1.64	0.0	3.7
5	0	0	0	0	1	0	0	0	0	0	0	1	1.51	1.51	0.0	3.7
6	1	0	0	0	1	0	0	0	0	0	0	2	1.52	1.83	0.32	11.1
7	0	0	0	0	1	0	0	0	0	0	0	1	1.57	1.57	0.0	3.7
8	0	0	0	0	1	0	0	0	0	0	0	1	1.51	1.51	0.0	3.7
9	1	0	0	0	1	0	0	0	0	0	0	2	1.24	1.42	0.18	11.1
10	0	0	0	0	1	0	0	0	0	0	0	1	1.6	1.6	0.0	3.7

10.2.1 Bar graph : Dihedral violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right


10.3 Dihedral-angle violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Number of violated restraints												Fraction of the ensemble	
NU0	ALPHA	GAMMA	NU2	CHI	ZETA	EPSILON	NU1	NU3	BETA	NU4	Total	Count ¹	%
2	0	0	0	0	0	0	0	0	0	0	2	1	10
0	0	0	0	0	0	0	0	0	0	0	0	2	20
0	0	0	0	0	0	0	0	0	0	0	0	3	30
0	0	0	0	0	0	0	0	0	0	0	0	4	40
0	0	0	0	0	0	0	0	0	0	0	0	5	50
0	0	0	0	0	0	0	0	0	0	0	0	6	60
0	0	0	0	0	0	0	0	0	0	0	0	7	70
0	0	0	0	0	0	0	0	0	0	0	0	8	80
0	0	0	0	0	0	0	0	0	0	0	0	9	90
0	0	0	0	1	0	0	0	0	0	0	1	10	100

¹ Number of models with violations


10.3.1 Bar graph : Dihedral-angle Violation statistics for the ensemble [i](#)



10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

10.4.1 Histogram : Distribution of mean dihedral-angle violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints [i](#)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.


Key	Atom-1	Atom-2	Atom-3	Atom-4	Models ¹	Mean	SD ²	Median
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	10	1.58	0.12	1.54

¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints [i](#)

10.5.1 Histogram : Distribution of violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints [i](#)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	6	1.83
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	2	1.72
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	4	1.64
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	10	1.6
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	7	1.57
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	3	1.52
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	5	1.51
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	8	1.51
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	1	1.44
(1,222)	1:7:A:A:O4'	1:7:A:A:C1'	1:7:A:A:N9	1:7:A:A:C4	9	1.42