



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 10:45 PM EDT

PDB ID : 5ZO7  
Title : Kinesin spindle protein Eg5 in complex with STLC-type inhibitor PVEI0138  
Authors : Yokoyama, H.; Sato, K.  
Deposited on : 2018-04-12  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	367	 4% 82% 8% 10%
1	B	367	 8% 79% 11% 10%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5441 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kinesin-like protein KIF11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	332	2621	1641	456	513	11	0	0	0
1	B	332	2621	1641	456	513	11	0	0	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	10	MET	-	expression tag	UNP P52732
A	11	ASN	-	expression tag	UNP P52732
A	12	HIS	-	expression tag	UNP P52732
A	13	LYS	-	expression tag	UNP P52732
A	14	VAL	-	expression tag	UNP P52732
A	15	HIS	-	expression tag	UNP P52732
A	16	MET	-	expression tag	UNP P52732
A	370	GLN	-	expression tag	UNP P52732
A	371	HIS	-	expression tag	UNP P52732
A	372	HIS	-	expression tag	UNP P52732
A	373	HIS	-	expression tag	UNP P52732
A	374	HIS	-	expression tag	UNP P52732
A	375	HIS	-	expression tag	UNP P52732
A	376	HIS	-	expression tag	UNP P52732
B	10	MET	-	expression tag	UNP P52732
B	11	ASN	-	expression tag	UNP P52732
B	12	HIS	-	expression tag	UNP P52732
B	13	LYS	-	expression tag	UNP P52732
B	14	VAL	-	expression tag	UNP P52732
B	15	HIS	-	expression tag	UNP P52732
B	16	MET	-	expression tag	UNP P52732
B	370	GLN	-	expression tag	UNP P52732
B	371	HIS	-	expression tag	UNP P52732
B	372	HIS	-	expression tag	UNP P52732
B	373	HIS	-	expression tag	UNP P52732

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	374	HIS	-	expression tag	UNP P52732
B	375	HIS	-	expression tag	UNP P52732
B	376	HIS	-	expression tag	UNP P52732

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	27	10	5	10	2	0	0
2	B	1	27	10	5	10	2	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	A	1	1	1	0	0
3	B	1	1	1	0	0

- Molecule 4 is (2R)-2-azanyl-3-[[2-(4-methoxyphenyl)-2-tricyclo[9.4.0.0<sup>3,8</sup>]pentadeca-1(11),3,5,7,12,14-hexaenyl]sulfanyl]propanoic acid (three-letter code: 5C5) (formula:  $C_{25}H_{25}NO_3S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			30	25	1	3	1		
4	B	1	Total	C	N	O	S	0	0
			30	25	1	3	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	A	47	Total 47	O 47	0	0
6	B	26	Total 26	O 26	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kinesin-like protein KIF11



- Molecule 1: Kinesin-like protein KIF11



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	158.35Å 50.58Å 93.69Å 90.00° 102.20° 90.00°	Depositor
Resolution (Å)	19.52 – 2.60 19.52 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.0 (19.52-2.60) 99.3 (19.52-2.60)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.39 (at 2.59Å)	Xtrriage
Refinement program	REFMAC 5.8.0048	Depositor
R, $R_{free}$	0.225 , 0.283 0.230 , 0.286	Depositor DCC
$R_{free}$ test set	2265 reflections (10.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.6	Xtrriage
Anisotropy	0.230	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 53.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5441	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0142e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5C5, SO4, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/2659	0.74	1/3594 (0.0%)
1	B	0.53	0/2659	0.72	0/3594
All	All	0.55	0/5318	0.73	1/7188 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	234	ARG	NE-CZ-NH2	-5.18	117.71	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2621	0	2645	12	0
1	B	2621	0	2645	26	0
2	A	27	0	12	0	0
2	B	27	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	30	0	0	0	0
4	B	30	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	5	0	0	0	0
5	B	5	0	0	1	0
6	A	47	0	0	0	0
6	B	26	0	0	0	0
All	All	5441	0	5314	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:242:THR:CG2	1:B:258:ILE:HD11	2.18	0.73
1:A:102:PHE:HB3	1:A:264:VAL:HB	1.75	0.69
1:B:102:PHE:HB3	1:B:264:VAL:HB	1.75	0.67
1:B:104:TYR:CE1	1:B:352:TYR:CE1	2.89	0.61
1:B:172:LEU:O	1:B:174:PRO:HD3	2.05	0.57
1:B:185:PHE:HE2	1:B:197:LYS:HG2	1.70	0.56
1:A:104:TYR:CE1	1:A:352:TYR:CE1	2.94	0.56
1:B:242:THR:HG22	1:B:258:ILE:HD11	1.86	0.56
1:B:102:PHE:CE1	1:B:332:ILE:HG12	2.42	0.54
1:B:185:PHE:CE2	1:B:197:LYS:HG2	2.44	0.53
1:B:82:TYR:OH	1:B:142:GLN:HG3	2.10	0.52
1:A:82:TYR:OH	1:A:142:GLN:HG3	2.11	0.51
1:A:102:PHE:CE1	1:A:332:ILE:HG12	2.46	0.49
1:B:144:PHE:O	1:B:148:THR:HB	2.13	0.48
1:B:323:SER:HA	1:B:328:THR:HB	1.96	0.48
1:A:234:ARG:HD3	1:A:288:ILE:CG2	2.44	0.47
1:B:33:ARG:NH1	5:B:603:SO4:O2	2.47	0.47
1:A:323:SER:HA	1:A:328:THR:HB	1.95	0.47
1:B:160:LEU:HB3	1:B:172:LEU:HG	1.96	0.47
1:A:160:LEU:HB3	1:A:172:LEU:HG	1.97	0.47
1:B:244:HIS:CE1	1:B:258:ILE:HD13	2.50	0.46
1:B:51:SER:HB3	1:B:63:ARG:NH1	2.31	0.46
1:B:185:PHE:HE2	1:B:197:LYS:HE3	1.81	0.46
1:B:29:ASN:OD1	1:B:29:ASN:C	2.55	0.45
1:A:29:ASN:OD1	1:A:29:ASN:C	2.56	0.45
1:B:185:PHE:CE2	1:B:197:LYS:HE3	2.53	0.44
1:B:152:THR:HG21	1:B:245:MET:HG2	2.00	0.43
1:B:172:LEU:O	1:B:174:PRO:CD	2.67	0.43
1:A:152:THR:HG21	1:A:245:MET:HG2	2.01	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:104:TYR:CD2	1:B:104:TYR:C	2.92	0.43
4:B:602:5C5:C19	4:B:602:5C5:C2	2.97	0.42
1:B:327:ARG:HA	1:B:362:LYS:O	2.20	0.42
1:A:104:TYR:CD2	1:A:104:TYR:C	2.94	0.41
1:B:18:ASN:ND2	1:B:360:LEU:HD23	2.35	0.41
1:A:43:CYS:O	1:A:45:PRO:HD3	2.21	0.41
1:B:23:VAL:HG11	1:B:50:VAL:HG21	2.03	0.41
1:B:241:VAL:CG1	1:B:261:LEU:HB3	2.51	0.41
1:B:258:ILE:CG2	1:B:366:ASN:O	2.68	0.41
1:A:106:GLN:O	1:A:109:THR:OG1	2.30	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/367 (89%)	314 (96%)	12 (4%)	0	100	100
1	B	326/367 (89%)	314 (96%)	12 (4%)	0	100	100
All	All	652/734 (89%)	628 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/324 (92%)	288 (97%)	10 (3%)	37	63
1	B	298/324 (92%)	288 (97%)	10 (3%)	37	63
All	All	596/648 (92%)	576 (97%)	20 (3%)	37	63

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	HIS
1	A	51	SER
1	A	87	CYS
1	A	175	SER
1	A	177	ASP
1	A	246	LYS
1	A	248	THR
1	A	251	ASP
1	A	291	SER
1	A	293	LEU
1	B	38	HIS
1	B	51	SER
1	B	87	CYS
1	B	248	THR
1	B	287	ASN
1	B	291	SER
1	B	293	LEU
1	B	305	ARG
1	B	329	ARG
1	B	362	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
1	A	20	GLN
1	A	183	GLN
1	A	287	ASN
1	B	18	ASN
1	B	287	ASN
1	B	366	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	5C5	B	602	-	32,33,33	2.06	4 (12%)	38,47,47	1.09	3 (7%)
5	SO4	A	603	-	4,4,4	0.33	0	6,6,6	1.18	0
2	ADP	B	600	3	24,29,29	0.81	1 (4%)	29,45,45	1.40	4 (13%)
5	SO4	B	603	-	4,4,4	0.39	0	6,6,6	0.27	0
2	ADP	A	600	3	24,29,29	1.10	2 (8%)	29,45,45	1.66	7 (24%)
4	5C5	A	602	-	32,33,33	2.11	4 (12%)	38,47,47	1.21	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	5C5	B	602	-	-	5/15/35/35	0/4/4/4
4	5C5	A	602	-	-	6/15/35/35	0/4/4/4
2	ADP	B	600	3	-	0/12/32/32	0/3/3/3

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	600	3	-	2/12/32/32	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	602	5C5	C9-C4	7.73	1.50	1.40
4	A	602	5C5	C12-C17	7.03	1.49	1.40
4	B	602	5C5	C9-C4	6.66	1.49	1.40
4	B	602	5C5	C12-C17	6.43	1.48	1.40
4	B	602	5C5	C3-S	-5.84	1.78	1.86
4	A	602	5C5	C3-S	-3.13	1.82	1.86
2	A	600	ADP	C2-N3	2.60	1.36	1.32
4	A	602	5C5	C3-C18	-2.57	1.50	1.54
2	A	600	ADP	C5-C4	2.48	1.47	1.40
4	B	602	5C5	C3-C17	-2.29	1.49	1.53
2	B	600	ADP	C5-C4	2.07	1.46	1.40

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	ADP	N3-C2-N1	-4.41	121.79	128.68
2	B	600	ADP	N3-C2-N1	-4.31	121.94	128.68
4	B	602	5C5	C18-C3-C17	3.90	119.13	111.81
4	A	602	5C5	C18-C3-C17	3.64	118.64	111.81
2	A	600	ADP	N6-C6-N1	3.52	125.88	118.57
2	A	600	ADP	PA-O3A-PB	-3.12	122.14	132.83
2	B	600	ADP	C2-N1-C6	2.98	123.84	118.75
2	A	600	ADP	C5-C6-N6	-2.92	115.92	120.35
2	A	600	ADP	O3'-C3'-C2'	-2.65	103.24	111.82
4	B	602	5C5	C17-C3-C4	-2.53	100.65	109.75
4	A	602	5C5	C22-O1-C21	2.51	122.95	117.51
4	A	602	5C5	C17-C3-C4	-2.47	100.86	109.75
2	A	600	ADP	C2-N1-C6	2.46	122.96	118.75
2	B	600	ADP	PA-O3A-PB	-2.36	124.73	132.83
2	B	600	ADP	N6-C6-N1	2.29	123.33	118.57
2	A	600	ADP	C3'-C2'-C1'	2.19	104.27	100.98
4	A	602	5C5	O-C-O2	-2.04	119.45	124.09
4	B	602	5C5	C18-C3-C4	2.03	115.62	111.81

There are no chirality outliers.

All (13) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	A	602	5C5	C24-C18-C3-S
4	A	602	5C5	C19-C18-C3-S
4	B	602	5C5	C24-C18-C3-S
4	B	602	5C5	C19-C18-C3-S
4	B	602	5C5	C23-C21-O1-C22
4	A	602	5C5	C23-C21-O1-C22
4	B	602	5C5	C20-C21-O1-C22
4	A	602	5C5	C20-C21-O1-C22
2	A	600	ADP	PA-O3A-PB-O2B
4	A	602	5C5	C24-C18-C3-C17
4	A	602	5C5	C19-C18-C3-C17
4	B	602	5C5	C24-C18-C3-C17
2	A	600	ADP	O4'-C4'-C5'-O5'



There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	602	5C5	1	0
5	B	603	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	332/367 (90%)	-0.00	15 (4%) 33 26	21, 41, 80, 114	0
1	B	332/367 (90%)	0.20	28 (8%) 11 7	31, 51, 88, 122	0
All	All	664/734 (90%)	0.10	43 (6%) 18 14	21, 46, 83, 122	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	122	ASN	4.8
1	B	59	ASP	4.7
1	A	38	HIS	4.5
1	B	149	ASP	4.4
1	B	38	HIS	4.3
1	A	151	GLY	4.0
1	B	208	ASP	3.9
1	B	122	ASN	3.9
1	B	150	ASN	3.6
1	A	305	ARG	3.6
1	B	121	PRO	3.6
1	B	308	HIS	3.5
1	B	189	ARG	3.5
1	B	124	GLU	3.5
1	B	61	SER	3.5
1	A	124	GLU	3.2
1	A	46	VAL	3.1
1	A	149	ASP	3.1
1	B	305	ARG	3.0
1	A	150	ASN	3.0
1	B	148	THR	3.0
1	B	254	GLU	2.9
1	A	53	ARG	2.8
1	B	46	VAL	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	16	MET	2.6
1	A	152	THR	2.6
1	B	185	PHE	2.6
1	B	125	TYR	2.6
1	B	151	GLY	2.6
1	B	153	GLU	2.4
1	A	308	HIS	2.4
1	B	60	LYS	2.4
1	B	145	GLU	2.4
1	B	197	LYS	2.2
1	B	23	VAL	2.1
1	B	190	ASN	2.1
1	A	23	VAL	2.1
1	A	123	GLU	2.1
1	B	152	THR	2.1
1	B	212	GLN	2.0
1	B	198	GLY	2.0
1	A	47	ARG	2.0
1	B	137	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	5C5	A	602	30/30	0.90	0.16	32,40,45,46	0
4	5C5	B	602	30/30	0.90	0.18	46,51,57,59	0
3	MG	A	601	1/1	0.93	0.16	36,36,36,36	0
3	MG	B	601	1/1	0.94	0.08	37,37,37,37	0

*Continued on next page...*

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ADP	A	600	27/27	0.96	0.15	30,33,37,38	0
2	ADP	B	600	27/27	0.96	0.16	35,39,42,43	0
5	SO4	A	603	5/5	0.98	0.14	40,42,43,45	0
5	SO4	B	603	5/5	0.98	0.10	48,51,52,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around 5C5 B 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ADP A 600:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ADP B 600:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.