



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2024 – 06:11 am BST

PDB ID : 4ZQH
Title : Crystal structure of DOX-P Reductoisomerase in complex with NADPH, fosmidomycin and magnesium
Authors : Birkinshaw, R.W.; Brady, R.L.
Deposited on : 2015-05-10
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	 85% 8% 6%
1	B	432	 84% 9% 6%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6450 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-deoxy-D-xylulose 5-phosphate reductoisomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	408	Total	C	N	O	S	0	1	0
			3055	1941	520	578	16			
1	B	408	Total	C	N	O	S	0	1	0
			3057	1943	520	578	16			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	initiating methionine	UNP A0A076U3E6
A	-16	ALA	-	expression tag	UNP A0A076U3E6
A	-15	HIS	-	expression tag	UNP A0A076U3E6
A	-14	HIS	-	expression tag	UNP A0A076U3E6
A	-13	HIS	-	expression tag	UNP A0A076U3E6
A	-12	HIS	-	expression tag	UNP A0A076U3E6
A	-11	HIS	-	expression tag	UNP A0A076U3E6
A	-10	HIS	-	expression tag	UNP A0A076U3E6
A	-9	SER	-	expression tag	UNP A0A076U3E6
A	-8	SER	-	expression tag	UNP A0A076U3E6
A	-7	GLY	-	expression tag	UNP A0A076U3E6
A	-6	LEU	-	expression tag	UNP A0A076U3E6
A	-5	GLU	-	expression tag	UNP A0A076U3E6
A	-4	VAL	-	expression tag	UNP A0A076U3E6
A	-3	LEU	-	expression tag	UNP A0A076U3E6
A	-2	PHE	-	expression tag	UNP A0A076U3E6
A	-1	GLN	-	expression tag	UNP A0A076U3E6
A	0	GLY	-	expression tag	UNP A0A076U3E6
A	1	PRO	-	expression tag	UNP A0A076U3E6
B	-17	MET	-	initiating methionine	UNP A0A076U3E6
B	-16	ALA	-	expression tag	UNP A0A076U3E6
B	-15	HIS	-	expression tag	UNP A0A076U3E6
B	-14	HIS	-	expression tag	UNP A0A076U3E6
B	-13	HIS	-	expression tag	UNP A0A076U3E6
B	-12	HIS	-	expression tag	UNP A0A076U3E6

Continued on next page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-11	HIS	-	expression tag	UNP A0A076U3E6
B	-10	HIS	-	expression tag	UNP A0A076U3E6
B	-9	SER	-	expression tag	UNP A0A076U3E6
B	-8	SER	-	expression tag	UNP A0A076U3E6
B	-7	GLY	-	expression tag	UNP A0A076U3E6
B	-6	LEU	-	expression tag	UNP A0A076U3E6
B	-5	GLU	-	expression tag	UNP A0A076U3E6
B	-4	VAL	-	expression tag	UNP A0A076U3E6
B	-3	LEU	-	expression tag	UNP A0A076U3E6
B	-2	PHE	-	expression tag	UNP A0A076U3E6
B	-1	GLN	-	expression tag	UNP A0A076U3E6
B	0	GLY	-	expression tag	UNP A0A076U3E6
B	1	PRO	-	expression tag	UNP A0A076U3E6

- # NDP

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is 3-[FORMYL(HYDROXY)AMINO]PROPYLPHOSPHONIC ACID (three-letter code: FOM) (formula: $C_4H_{10}NO_5P$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P 11 4 1 5 1	0	0
4	B	1	Total C N O P 11 4 1 5 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is 2-[N-CYCLOHEXYLAMINO]ETHANE SULFONIC ACID (three-letter code: NHE) (formula: C₈H₁₇NO₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			13	8	1	3	1		
6	B	1	Total	C	N	O	S	0	0
			13	8	1	3	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	82	Total	O	0	0
			82	82		
7	B	80	Total	O	0	0
			80	80		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1-deoxy-D-xylulose 5-phosphate reductoisomerase

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	65.72Å 65.72Å 391.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	65.24 – 2.40 65.24 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (65.24-2.40) 100.0 (65.24-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.40Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.202 , 0.243 0.203 , 0.242	Depositor DCC
R_{free} test set	1856 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	36.7	Xtriage
Anisotropy	0.580	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.489 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6450	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FOM, NDP, NHE, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/3102	0.67	0/4208
1	B	0.54	0/3104	0.67	0/4210
All	All	0.55	0/6206	0.67	0/8418

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3055	0	3108	21	0
1	B	3057	0	3112	24	0
2	A	48	0	26	0	0
2	B	48	0	26	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	11	0	7	0	0
4	B	11	0	7	0	0
5	A	12	0	16	2	0
5	B	18	0	24	3	0
6	A	13	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	13	0	16	2	0
7	A	82	0	0	1	0
7	B	80	0	0	3	0
All	All	6450	0	6358	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:SER:O	1:A:73:THR:HG23	1.90	0.72
1:B:69:SER:O	1:B:73:THR:HG23	1.90	0.72
1:A:155:GLU:OE1	5:A:504:GOL:H31	1.91	0.70
1:B:257:SER:HA	5:B:505:GOL:H12	1.77	0.67
1:B:224:ASP:OD2	7:B:601:HOH:O	2.15	0.65
1:A:38:THR:OG1	7:A:601:HOH:O	2.14	0.64
1:A:181:ILE:HD13	1:A:265:TYR:CE2	2.33	0.64
1:B:181:ILE:HD13	1:B:265:TYR:CE2	2.33	0.64
1:A:300:LEU:HD11	1:A:305:LEU:HD21	1.82	0.61
1:B:300:LEU:HD11	1:B:305:LEU:HD21	1.81	0.61
1:A:155:GLU:OE1	5:A:504:GOL:C3	2.52	0.57
1:B:326:TYR:CE1	1:B:330:LEU:HD11	2.39	0.57
1:A:326:TYR:CE1	1:A:330:LEU:HD11	2.40	0.57
1:B:114:LEU:HD23	1:B:387:LEU:HD22	1.89	0.53
1:B:257:SER:HA	5:B:505:GOL:C1	2.41	0.51
1:B:243:ASP:O	1:B:243:ASP:OD2	2.29	0.50
1:A:243:ASP:O	1:A:243:ASP:OD2	2.29	0.49
1:B:57:LYS:NZ	1:B:94:ASP:OD2	2.35	0.49
1:B:58:ILE:HD12	1:B:68:LEU:HD21	1.95	0.49
1:A:58:ILE:HD12	1:A:68:LEU:HD21	1.95	0.48
6:B:507:NHE:H6'1	7:B:617:HOH:O	2.14	0.47
1:A:307:SER:HB2	1:B:307:SER:HB2	1.97	0.46
1:B:260:HIS:O	1:B:261:SER:CB	2.64	0.45
1:B:240:HIS:CE1	6:B:507:NHE:H6'2	2.51	0.45
1:A:260:HIS:O	1:A:261:SER:CB	2.65	0.44
1:B:29:PRO:O	1:B:30:ASN:HB2	2.17	0.44
1:A:393:MET:O	1:A:397:VAL:HG23	2.18	0.44
1:A:57:LYS:NZ	1:A:94:ASP:OD2	2.35	0.44
1:B:393:MET:O	1:B:397:VAL:HG23	2.18	0.44
1:A:22:LEU:HD23	1:A:22:LEU:HA	1.89	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:134:ALA:HB2	1:A:390:ILE:HD13	1.99	0.44
1:B:134:ALA:HB2	1:B:390:ILE:HD13	1.99	0.43
1:B:16:SER:OG	5:B:506:GOL:C3	2.67	0.43
1:A:144:LYS:HD2	1:A:145:HIS:CE1	2.54	0.43
1:B:181:ILE:CD1	1:B:265:TYR:CE2	3.00	0.43
1:B:247:HIS:O	1:B:250:GLN:NE2	2.52	0.43
1:A:247:HIS:O	1:A:250:GLN:NE2	2.51	0.42
1:A:77:LEU:HB3	1:A:79:THR:HG22	2.02	0.42
1:B:377:SER:CB	7:B:667:HOH:O	2.68	0.42
1:B:77:LEU:HB3	1:B:79:THR:HG22	2.02	0.41
1:A:181:ILE:CD1	1:A:265:TYR:CE2	3.00	0.41
1:A:134:ALA:O	1:A:138:VAL:HG23	2.21	0.41
1:B:359:ILE:O	1:B:363:VAL:HG23	2.21	0.41
1:A:359:ILE:O	1:A:363:VAL:HG23	2.20	0.41
1:B:134:ALA:O	1:B:138:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	407/432 (94%)	393 (97%)	12 (3%)	2 (0%)	29	41
1	B	407/432 (94%)	393 (97%)	12 (3%)	2 (0%)	29	41
All	All	814/864 (94%)	786 (97%)	24 (3%)	4 (0%)	29	41

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	SER
1	B	261	SER
1	B	191	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	191	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/349 (92%)	311 (97%)	9 (3%)	43	63
1	B	320/349 (92%)	313 (98%)	7 (2%)	52	71
All	All	640/698 (92%)	624 (98%)	16 (2%)	47	67

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	THR
1	A	67	GLN
1	A	79	THR
1	A	114	LEU
1	A	144	LYS
1	A	228	MET
1	A	231	LYS
1	A	262	LEU
1	A	371	THR
1	B	38	THR
1	B	67	GLN
1	B	79	THR
1	B	228	MET
1	B	231	LYS
1	B	262	LEU
1	B	371	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	30	ASN
1	A	67	GLN
1	A	145	HIS
1	A	247	HIS
1	A	304	GLN
1	A	374	GLN
1	B	27	GLN
1	B	64	ASN
1	B	240	HIS
1	B	247	HIS
1	B	304	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	B	502	-	45,52,52	1.04	2 (4%)	53,80,80	1.20	5 (9%)
4	FOM	A	503	3	9,10,10	2.22	3 (33%)	11,13,13	2.57	4 (36%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FOM	B	504	3	9,10,10	2.98	5 (55%)	11,13,13	2.44	5 (45%)
5	GOL	B	501	-	5,5,5	0.30	0	5,5,5	0.53	0
5	GOL	B	506	-	5,5,5	0.18	0	5,5,5	0.73	0
5	GOL	A	506	-	5,5,5	0.42	0	5,5,5	0.74	0
2	NDP	A	501	-	45,52,52	1.11	3 (6%)	53,80,80	1.19	4 (7%)
5	GOL	A	504	-	5,5,5	0.74	0	5,5,5	1.11	0
6	NHE	A	505	-	13,13,13	2.81	2 (15%)	16,17,17	1.68	3 (18%)
5	GOL	B	505	-	5,5,5	0.60	0	5,5,5	1.15	1 (20%)
6	NHE	B	507	-	13,13,13	2.88	2 (15%)	16,17,17	1.87	3 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	B	502	-	-	3/30/77/77	0/5/5/5
4	FOM	A	503	3	-	0/7/9/9	-
4	FOM	B	504	3	-	0/7/9/9	-
5	GOL	B	501	-	-	0/4/4/4	-
5	GOL	B	506	-	-	2/4/4/4	-
5	GOL	A	506	-	-	2/4/4/4	-
2	NDP	A	501	-	-	5/30/77/77	0/5/5/5
5	GOL	A	504	-	-	2/4/4/4	-
6	NHE	A	505	-	-	4/7/15/15	0/1/1/1
5	GOL	B	505	-	-	4/4/4/4	-
6	NHE	B	507	-	-	6/7/15/15	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	507	NHE	C2-S	-7.67	1.66	1.77
6	A	505	NHE	C2-S	-7.61	1.66	1.77
6	B	507	NHE	O2-S	6.84	1.65	1.45
6	A	505	NHE	O1-S	6.43	1.64	1.45
4	B	504	FOM	PA1-OP1	5.09	1.60	1.50
4	A	503	FOM	C1-N1	-4.83	1.27	1.34
4	B	504	FOM	C1-N1	-4.78	1.27	1.34
4	B	504	FOM	PA1-OP2	-3.77	1.46	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	NDP	C6N-C5N	3.49	1.39	1.33
2	B	502	NDP	C6N-C5N	3.27	1.39	1.33
4	B	504	FOM	PA1-C4	3.22	1.82	1.78
4	A	503	FOM	PA1-OP3	3.02	1.61	1.54
2	A	501	NDP	O4B-C1B	2.73	1.44	1.41
2	B	502	NDP	O4B-C1B	2.65	1.44	1.41
4	B	504	FOM	PA1-OP3	2.49	1.60	1.54
4	A	503	FOM	PA1-C4	2.44	1.81	1.78
2	A	501	NDP	P2B-O2B	2.40	1.63	1.59

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	FOM	PA1-C4-C3	-5.23	108.44	114.98
6	B	507	NHE	C1-N-C1'	4.99	123.94	114.14
4	A	503	FOM	C3-C2-N1	-4.87	101.06	111.07
4	B	504	FOM	C3-C2-N1	-4.60	101.60	111.07
6	A	505	NHE	C1-N-C1'	4.54	123.05	114.14
4	B	504	FOM	PA1-C4-C3	-4.45	109.41	114.98
6	B	507	NHE	O3-S-C2	4.09	112.39	105.77
2	B	502	NDP	N3A-C2A-N1A	-3.99	122.45	128.68
6	A	505	NHE	O2-S-C2	3.30	110.88	106.92
2	A	501	NDP	N3A-C2A-N1A	-3.12	123.81	128.68
2	A	501	NDP	O3X-P2B-O2X	2.64	117.72	107.64
4	A	503	FOM	OP3-PA1-OP2	2.59	115.65	108.08
2	B	502	NDP	PN-O3-PA	-2.46	124.40	132.83
4	B	504	FOM	OP2-PA1-OP1	-2.43	105.96	112.39
2	B	502	NDP	N6A-C6A-N1A	2.42	123.60	118.57
2	A	501	NDP	O2N-PN-O1N	2.42	124.20	112.24
6	A	505	NHE	O3-S-C2	2.39	109.64	105.77
4	B	504	FOM	O1-C1-N1	-2.36	119.00	125.80
2	B	502	NDP	O3X-P2B-O2X	2.35	116.62	107.64
4	A	503	FOM	OP3-PA1-OP1	-2.35	106.18	112.39
2	A	501	NDP	O3D-C3D-C4D	-2.33	104.32	111.05
6	B	507	NHE	O1-S-C2	2.30	109.68	106.92
4	B	504	FOM	OP2-PA1-C4	2.26	112.27	106.95
2	B	502	NDP	C5A-C6A-N6A	-2.18	117.04	120.35
5	B	505	GOL	O2-C2-C3	2.02	118.03	109.12

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	NDP	C2B-O2B-P2B-O3X
5	A	504	GOL	O1-C1-C2-C3
5	A	506	GOL	C1-C2-C3-O3
5	B	505	GOL	O1-C1-C2-O2
5	B	505	GOL	C1-C2-C3-O3
5	B	506	GOL	O1-C1-C2-C3
6	A	505	NHE	C6'-C1'-N-C1
6	B	507	NHE	C6'-C1'-N-C1
6	B	507	NHE	N-C1-C2-S
6	B	507	NHE	C1-C2-S-O1
6	B	507	NHE	C1-C2-S-O2
6	B	507	NHE	C1-C2-S-O3
5	B	505	GOL	O1-C1-C2-C3
5	B	505	GOL	O2-C2-C3-O3
5	B	506	GOL	O1-C1-C2-O2
6	A	505	NHE	C1-C2-S-O3
6	B	507	NHE	C2'-C1'-N-C1
5	A	506	GOL	O2-C2-C3-O3
2	A	501	NDP	O4D-C1D-N1N-C6N
5	A	504	GOL	O1-C1-C2-O2
2	B	502	NDP	O4D-C1D-N1N-C6N
6	A	505	NHE	C1-C2-S-O2
2	A	501	NDP	O4B-C4B-C5B-O5B
2	A	501	NDP	C2B-O2B-P2B-O2X
2	A	501	NDP	C5D-O5D-PN-O3
2	B	502	NDP	C5D-O5D-PN-O3
2	B	502	NDP	O4B-C4B-C5B-O5B
6	A	505	NHE	C1-C2-S-O1


There are no ring outliers.


4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	506	GOL	1	0
5	A	504	GOL	2	0
5	B	505	GOL	2	0
6	B	507	NHE	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	408/432 (94%)	-0.51	0 100 100	39, 57, 81, 98	0
1	B	408/432 (94%)	-0.52	1 (0%) 95 94	39, 57, 80, 106	0
All	All	816/864 (94%)	-0.52	1 (0%) 95 95	39, 57, 81, 106	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	408	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NHE	B	507	13/13	0.79	0.21	77,87,105,107	0
6	NHE	A	505	13/13	0.82	0.21	80,90,105,115	0
5	GOL	A	506	6/6	0.92	0.13	66,68,71,78	0
5	GOL	A	504	6/6	0.93	0.11	53,61,63,65	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	B	501	6/6	0.96	0.12	65,68,71,76	0
5	GOL	B	505	6/6	0.96	0.16	56,60,61,68	0
5	GOL	B	506	6/6	0.97	0.12	52,58,59,64	0
4	FOM	A	503	11/11	0.98	0.10	41,47,49,50	0
2	NDP	A	501	48/48	0.98	0.10	42,48,53,58	0
2	NDP	B	502	48/48	0.98	0.11	42,48,55,58	0
3	MG	A	502	1/1	0.98	0.12	43,43,43,43	0
4	FOM	B	504	11/11	0.99	0.11	41,46,49,51	0
3	MG	B	503	1/1	0.99	0.10	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.