



wwPDB EM Validation Summary Report ⓘ

Apr 9, 2025 – 04:38 pm BST

PDB ID : 7ZU0 / pdb_00007zu0
EMDB ID : EMD-14964
Title : HOPS tethering complex from yeast
Authors : Shvarev, D.; Schoppe, J.; Koenig, C.; Perz, A.; Fuellbrunn, N.; Kiontke, S.;
Langemeyer, L.; Janulienė, D.; Schnelle, K.; Kuemmel, D.; Froehlich, F.;
Moeller, A.; Ungermann, C.
Deposited on : 2022-05-11
Resolution : 4.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev117
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1029	
2	B	798	
3	C	918	
4	D	691	
5	E	1049	
6	F	1016	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 25763 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase PEP5.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	954	Total	C	N	O	S	0	0
			6914	4395	1185	1315	19		

- Molecule 2 is a protein called Vacuolar protein sorting-associated protein 16.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	733	Total	C	N	O	S	0	0
			5981	3862	980	1110	29		

- Molecule 3 is a protein called Vacuolar membrane protein PEP3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	842	Total	C	N	O	S	0	0
			5549	3476	989	1063	21		

- Molecule 4 is a protein called Vacuolar protein sorting-associated protein 33.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	597	Total	C	N	O	S	0	0
			4749	3036	789	904	20		

- Molecule 5 is a protein called Vacuolar morphogenesis protein 6.


Mol	Chain	Residues	Atoms				AltConf	Trace
5	E	343	Total	C	N	O	0	0
			1704	1018	343	343		

- Molecule 6 is a protein called Vacuolar protein sorting-associated protein 41.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	112	Total	C	N	O	S	0	0
			866	549	144	170	3		


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	993	ASP	-	expression tag	UNP P38959
F	994	TYR	-	expression tag	UNP P38959
F	995	LYS	-	expression tag	UNP P38959
F	996	ASP	-	expression tag	UNP P38959
F	997	ASP	-	expression tag	UNP P38959
F	998	ASP	-	expression tag	UNP P38959
F	999	ASP	-	expression tag	UNP P38959
F	1000	LYS	-	expression tag	UNP P38959
F	1001	ASP	-	expression tag	UNP P38959
F	1002	TYR	-	expression tag	UNP P38959
F	1003	LYS	-	expression tag	UNP P38959
F	1004	ASP	-	expression tag	UNP P38959
F	1005	ASP	-	expression tag	UNP P38959
F	1006	ASP	-	expression tag	UNP P38959
F	1007	ASP	-	expression tag	UNP P38959
F	1008	LYS	-	expression tag	UNP P38959
F	1009	ASP	-	expression tag	UNP P38959
F	1010	TYR	-	expression tag	UNP P38959
F	1011	LYS	-	expression tag	UNP P38959
F	1012	ASP	-	expression tag	UNP P38959
F	1013	ASP	-	expression tag	UNP P38959
F	1014	ASP	-	expression tag	UNP P38959
F	1015	ASP	-	expression tag	UNP P38959
F	1016	LYS	-	expression tag	UNP P38959




• Molecule 2: Vacuolar protein sorting-associated protein 16


Chain B: 32% 66% 25% 8%




• Molecule 3: Vacuolar membrane protein PEP3



• Molecule 4: Vacuolar protein sorting-associated protein 33





[illegible]

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	244661	Depositor
Resolution determination method	OTHER	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2800	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.893	Depositor
Minimum map value	-0.818	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.027	Depositor
Recommended contour level	0.35	Depositor
Map size (\AA)	620.92804, 620.92804, 620.92804	wwPDB
Map dimensions	672, 672, 672	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.9240001, 0.9240001, 0.9240001	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/7025	0.50	9/9555 (0.1%)
2	B	0.28	0/6101	0.49	1/8240 (0.0%)
3	C	0.26	0/5601	0.49	9/7644 (0.1%)
4	D	0.26	0/4827	0.47	2/6533 (0.0%)
5	E	0.44	0/1702	0.65	12/2373 (0.5%)
6	F	0.28	0/875	0.52	1/1180 (0.1%)
All	All	0.28	0/26131	0.50	34/35525 (0.1%)

There are no bond length outliers.

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	1002	PRO	N-CA-CB	6.80	111.46	103.30
3	C	134	PRO	N-CA-CB	6.75	111.40	103.30
3	C	216	PRO	N-CA-CB	6.63	111.26	103.30
3	C	195	PRO	N-CA-CB	6.58	111.20	103.30
5	E	900	PRO	N-CA-CB	6.50	111.10	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6914	0	6244	166	0
2	B	5981	0	6031	148	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	5549	0	4384	129	0
4	D	4749	0	4710	96	0
5	E	1704	0	731	20	0
6	F	866	0	830	23	0
All	All	25763	0	22930	546	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 546 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:479:ARG:HH22	1:A:490:ASP:HA	1.47	0.79
2:B:153:LEU:HB2	2:B:157:HIS:CE1	2.19	0.78
3:C:834:GLN:HA	3:C:838:PHE:HZ	1.48	0.77
2:B:447:LYS:HG2	2:B:486:VAL:HG21	1.67	0.76
1:A:750:SER:HB2	3:C:627:ILE:HD13	1.65	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	940/1029 (91%)	840 (89%)	97 (10%)	3 (0%)	37	72
2	B	729/798 (91%)	646 (89%)	82 (11%)	1 (0%)	48	83
3	C	820/918 (89%)	751 (92%)	61 (7%)	8 (1%)	13	48
4	D	583/691 (84%)	548 (94%)	35 (6%)	0	100	100
5	E	339/1049 (32%)	320 (94%)	14 (4%)	5 (2%)	8	39
6	F	108/1016 (11%)	96 (89%)	12 (11%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	3519/5501 (64%)	3201 (91%)	301 (9%)	17 (0%)	27	63

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	938	PRO
2	B	27	PRO
3	C	130	PRO
3	C	134	PRO
3	C	195	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	652/952 (68%)	652 (100%)	0	100	100
2	B	666/730 (91%)	666 (100%)	0	100	100
3	C	396/873 (45%)	396 (100%)	0	100	100
4	D	526/640 (82%)	526 (100%)	0	100	100
6	F	92/938 (10%)	92 (100%)	0	100	100
All	All	2332/4133 (56%)	2332 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	333	ASN
2	B	494	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	C	10
5	E	1
6	F	1
1	A	1

The worst 5 of 13 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	115:TYR	C	121:TRP	N	13.77
1	E	991:HIS	C	998:ASN	N	13.23
1	C	23:GLN	C	27:LEU	N	12.99
1	C	140:PHE	C	150:GLU	N	12.97
1	C	11:LEU	C	17:LYS	N	10.46


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-14964. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 336



Y Index: 336




Z Index: 336


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 342



Y Index: 372



Z Index: 358


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.35. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 164 nm^3 ; this corresponds to an approximate mass of 149 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.227 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-14964 and PDB model 7ZU0. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.35 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.35).

9.4 Atom inclusion [i](#)



At the recommended contour level, 74% of all backbone atoms, 60% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.35) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.5980	<div></div> 0.3930
A	<div></div> 0.5650	<div></div> 0.3590
B	<div></div> 0.5310	<div></div> 0.4490
C	<div></div> 0.8000	<div></div> 0.3510
D	<div></div> 0.4710	<div></div> 0.4620
E	<div></div> 0.6740	<div></div> 0.2820
F	<div></div> 0.5500	<div></div> 0.4010

1.0

0.0

<0.0