



## wwPDB EM Validation Summary Report ⓘ

Apr 23, 2024 – 10:28 am BST

PDB ID : 6ZVJ  
EMDB ID : EMD-11458  
Title : Structure of a human ABCE1-bound 43S pre-initiation complex - State II  
Authors : Kratzat, H.; Mackens-Kiani, T.; Ameismeier, A.; Cheng, J.; Berninghausen, O.; Becker, T.; Beckmann, R.  
Deposited on : 2020-07-24  
Resolution : 3.80 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92  
Mogul : 1.8.4, CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	a	216	 97%
2	p	211	 99%
3	d	216	 97%
4	Q	101	 100%
5	q	255	 98%
6	W	24	 100%
7	r	222	 99%
8	s	181	 94%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	t	205	 97%
10	c	180	 99%
11	n	144	 93% 6%
12	m	149	 99%
13	i	125	 100%
14	y	82	 98%
15	f	129	 98%
16	j	139	 98%
17	z	122	 98%
18	R	82	 99%
19	T	44	 98%
20	2	1720	 48% 42% 9%
21	w	130	 95%
22	g	138	 98%
23	b	224	 99%
24	e	189	 96%
25	u	95	 98%
26	v	116	 95%
27	o	119	 98%
28	k	140	 98%
29	x	141	 98%
30	h	98	 99%
31	P	70	 99%
32	S	61	 98%
33	l	54	 96%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
34	U	62	
35	V	309	
36	I	316	
37	B	566	
38	A	722	
39	C	743	
40	E	429	
41	F	269	
42	H	318	
43	K	217	
44	L	372	
45	M	362	
46	1	595	
47	N	91	
48	D	527	
49	Y	78	
50	G	258	
50	J	258	

## 2 Entry composition

There are 55 unique types of molecules in this entry. The entry contains 115020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 40S ribosomal protein SA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	a	216	Total	C	N	O	S	0	0
			1706	1083	299	316	8		

- Molecule 2 is a protein called 40S ribosomal protein S3a.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	p	211	Total	C	N	O	S	0	0
			1716	1088	307	307	14		

- Molecule 3 is a protein called 40S ribosomal protein S2.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	d	216	Total	C	N	O	S	0	0
			1675	1085	287	293	10		

- Molecule 4 is a protein called 40S ribosomal protein S26.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	Q	101	Total	C	N	O	S	0	0
			815	507	170	133	5		

- Molecule 5 is a protein called 40S ribosomal protein S4, X isoform.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	q	255	Total	C	N	O	S	0	0
			2032	1299	377	348	8		

- Molecule 6 is a protein called 60S ribosomal protein L41.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	W	24	Total	C	N	O	S	0	0
			231	139	62	27	3		

- Molecule 7 is a protein called 40S ribosomal protein S6.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	r	222	Total	C	N	O	S	0	0
			1795	1123	357	309	6		

- Molecule 8 is a protein called 40S ribosomal protein S7.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	s	173	Total	C	N	O	S	0	0
			1400	898	256	245	1		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
s	8	ILE	GLY	conflict	UNP P62081
s	9	VAL	GLU	conflict	UNP P62081
s	12	ASN	ASP	conflict	UNP P62081

- Molecule 9 is a protein called 40S ribosomal protein S8.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	t	199	Total	C	N	O	S	0	0
			1639	1027	322	285	5		

- Molecule 10 is a protein called 40S ribosomal protein S9.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	c	180	Total	C	N	O	S	0	0
			1500	955	300	243	2		

- Molecule 11 is a protein called 40S ribosomal protein S11.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	n	135	Total	C	N	O	S	0	0
			1120	715	211	188	6		

- Molecule 12 is a protein called 40S ribosomal protein S13.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	m	149	Total	C	N	O	S	0	0
			1203	770	228	204	1		

- Molecule 13 is a protein called 40S ribosomal protein S14.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	i	125	Total	C	N	O	S	0	0
			940	574	187	173	6		

- Molecule 14 is a protein called 40S ribosomal protein S21.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	y	82	Total	C	N	O	S	0	0
			626	384	116	121	5		

- Molecule 15 is a protein called 40S ribosomal protein S15a.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	f	129	Total	C	N	O	S	0	0
			1034	659	193	176	6		

- Molecule 16 is a protein called 40S ribosomal protein S23.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	j	139	Total	C	N	O	S	0	0
			1081	682	214	182	3		

- Molecule 17 is a protein called 40S ribosomal protein S24.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	z	122	Total	C	N	O	S	0	0
			1000	633	196	166	5		

- Molecule 18 is a protein called 40S ribosomal protein S27.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	R	82	Total	C	N	O	S	0	0
			641	402	118	114	7		

- Molecule 19 is a protein called 40S ribosomal protein S30.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	T	44	Total	C	N	O	S	0	0
			355	216	81	57	1		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
T	54	GLY	VAL	conflict	UNP P62861
T	56	ASN	THR	conflict	UNP P62861
T	57	ALA	PHE	conflict	UNP P62861
T	58	ASN	GLY	conflict	UNP P62861

- Molecule 20 is a RNA chain called 18S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	2	1720	Total	C	N	O	P	0	0
			36700	16391	6600	11998	1711		

- Molecule 21 is a protein called 40S ribosomal protein S17.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	w	128	Total	C	N	O	S	0	0
			1011	641	182	184	4		

- Molecule 22 is a protein called 40S ribosomal protein S16.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	g	138	Total	C	N	O	S	0	0
			1100	699	208	190	3		

- Molecule 23 is a protein called 40S ribosomal protein S3.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	b	224	Total	C	N	O	S	0	0
			1746	1112	314	313	7		

- Molecule 24 is a protein called 40S ribosomal protein S5.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	e	189	Total	C	N	O	S	0	0
			1495	934	284	270	7		

- Molecule 25 is a protein called 40S ribosomal protein S10.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	u	95	Total	C	N	O	S	0	0
			800	524	139	131	6		

- Molecule 26 is a protein called 40S ribosomal protein S12.



Mol	Chain	Residues	Atoms					AltConf	Trace
26	v	111	Total	C	N	O	S	0	0
			862	544	151	160	7		

- Molecule 27 is a protein called 40S ribosomal protein S15.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	o	119	Total	C	N	O	S	0	0
			980	623	183	167	7		

- Molecule 28 is a protein called 40S ribosomal protein S18.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	k	140	Total	C	N	O	S	0	0
			1163	731	234	197	1		

- Molecule 29 is a protein called 40S ribosomal protein S19.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	x	141	Total	C	N	O	S	0	0
			1095	685	210	197	3		

- Molecule 30 is a protein called 40S ribosomal protein S20.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	h	98	Total	C	N	O	S	0	0
			781	489	148	140	4		

- Molecule 31 is a protein called 40S ribosomal protein S25.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	P	70	Total	C	N	O	S	0	0
			558	358	101	98	1		

- Molecule 32 is a protein called 40S ribosomal protein S28.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	S	61	Total	C	N	O	S	0	0
			480	292	95	91	2		

- Molecule 33 is a protein called 40S ribosomal protein S29.

Mol	Chain	Residues	Atoms					AltConf	Trace
33	1	54	Total	C	N	O	S	0	0
			451	282	93	71	5		

- Molecule 34 is a protein called Ubiquitin-40S ribosomal protein S27a.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	U	57	Total	C	N	O	S	0	0
			466	295	89	75	7		

- Molecule 35 is a protein called Receptor of activated protein C kinase 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
35	V	296	Total	C	N	O	S	0	0
			2315	1464	404	435	12		

- Molecule 36 is a protein called Eukaryotic translation initiation factor 3 subunit I.

Mol	Chain	Residues	Atoms					AltConf	Trace
36	I	305	Total	C	N	O	S	0	0
			1497	887	305	305			

- Molecule 37 is a protein called Eukaryotic translation initiation factor 3 subunit B.

Mol	Chain	Residues	Atoms					AltConf	Trace
37	B	536	Total	C	N	O	S	0	0
			2966	1801	580	580	5		

- Molecule 38 is a protein called Eukaryotic translation initiation factor 3 subunit A.

Mol	Chain	Residues	Atoms					AltConf	Trace
38	A	692	Total	C	N	O	S	0	0
			5384	3378	980	1004	22		

- Molecule 39 is a protein called Eukaryotic translation initiation factor 3 subunit C.

Mol	Chain	Residues	Atoms					AltConf	Trace
39	C	625	Total	C	N	O	S	0	0
			5073	3205	899	934	35		

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	130	MET	-	insertion	UNP B4DVU3
C	131	ASN	HIS	conflict	UNP B4DVU3
C	132	LYS	ASP	conflict	UNP B4DVU3
C	133	ASN	ARG	conflict	UNP B4DVU3
C	134	ASN	LYS	conflict	UNP B4DVU3
C	135	ALA	SER	conflict	UNP B4DVU3
C	137	ALA	ARG	conflict	UNP B4DVU3
C	?	-	ASP	deletion	UNP B4DVU3
C	?	-	GLU	deletion	UNP B4DVU3
C	?	-	GLU	deletion	UNP B4DVU3
C	?	-	GLU	deletion	UNP B4DVU3
C	?	-	GLU	deletion	UNP B4DVU3
C	?	-	ASP	deletion	UNP B4DVU3
C	?	-	ASN	deletion	UNP B4DVU3
C	?	-	GLU	deletion	UNP B4DVU3
C	?	-	GLY	deletion	UNP B4DVU3
C	?	-	GLY	deletion	UNP B4DVU3
C	?	-	GLU	deletion	UNP B4DVU3
C	?	-	TRP	deletion	UNP B4DVU3
C	?	-	GLU	deletion	UNP B4DVU3
C	?	-	ARG	deletion	UNP B4DVU3
C	?	-	VAL	deletion	UNP B4DVU3
C	?	-	ARG	deletion	UNP B4DVU3
C	?	-	GLY	deletion	UNP B4DVU3
C	?	-	GLY	deletion	UNP B4DVU3
C	?	-	VAL	deletion	UNP B4DVU3
C	139	SER	PRO	conflict	UNP B4DVU3
C	140	THR	LEU	conflict	UNP B4DVU3
C	141	LEU	VAL	conflict	UNP B4DVU3
C	142	ARG	LYS	conflict	UNP B4DVU3
C	143	GLN	GLU	conflict	UNP B4DVU3
C	145	ILE	PRO	conflict	UNP B4DVU3
C	146	ARG	LYS	conflict	UNP B4DVU3
C	147	LYS	MET	conflict	UNP B4DVU3
C	148	TYR	PHE	conflict	UNP B4DVU3
C	149	ASN	ALA	conflict	UNP B4DVU3

- Molecule 40 is a protein called Eukaryotic translation initiation factor 3 subunit E.

Mol	Chain	Residues	Atoms					AltConf	Trace
40	E	416	Total	C	N	O	S	0	0
			3437	2202	585	630	20		

- Molecule 41 is a protein called Eukaryotic translation initiation factor 3 subunit F.

Mol	Chain	Residues	Atoms					AltConf	Trace
41	F	269	Total	C	N	O	S	0	0
			2090	1317	356	405	12		

- Molecule 42 is a protein called Eukaryotic translation initiation factor 3 subunit H.

Mol	Chain	Residues	Atoms					AltConf	Trace
42	H	295	Total	C	N	O	S	0	0
			2413	1532	417	449	15		

- Molecule 43 is a protein called Eukaryotic translation initiation factor 3 subunit K.

Mol	Chain	Residues	Atoms					AltConf	Trace
43	K	217	Total	C	N	O	S	0	0
			1750	1116	288	334	12		

- Molecule 44 is a protein called Eukaryotic translation initiation factor 3 subunit L.

Mol	Chain	Residues	Atoms					AltConf	Trace
44	L	372	Total	C	N	O	S	0	0
			3112	2011	520	564	17		

- Molecule 45 is a protein called Eukaryotic translation initiation factor 3 subunit M.

Mol	Chain	Residues	Atoms					AltConf	Trace
45	M	340	Total	C	N	O	S	0	0
			2718	1734	459	508	17		

- Molecule 46 is a protein called ATP-binding cassette sub-family E member 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
46	1	576	Total	C	N	O	S	0	0
			4542	2899	779	836	28		

- Molecule 47 is a protein called Eukaryotic translation initiation factor 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
47	N	91	Total	C	N	O	S	0	0
			728	460	131	135	2		

- Molecule 48 is a protein called Eukaryotic translation initiation factor 3 subunit D.

Mol	Chain	Residues	Atoms					AltConf	Trace
48	D	447	Total	C	N	O	S	0	0
			3618	2279	625	692	22		

- Molecule 49 is a protein called RNA recognition motif (unknown).

Mol	Chain	Residues	Atoms				AltConf	Trace
49	Y	78	Total	C	N	O	0	0
			390	234	78	78		


- Molecule 50 is a protein called Eukaryotic translation initiation factor 3 subunit J.

Mol	Chain	Residues	Atoms				AltConf	Trace
50	G	70	Total	C	N	O	0	0
			348	208	70	70		
50	J	73	Total	C	N	O	0	0
			363	217	73	73		

- Molecule 51 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
51	Q	1	Total	Zn	0
			1	1	
51	l	1	Total	Zn	0
			1	1	
51	U	1	Total	Zn	0
			1	1	

- Molecule 52 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>).



Mol	Chain	Residues	Atoms			AltConf
52	1	1	Total	Fe	S	0
			8	4	4	
52	1	1	Total	Fe	S	0
			8	4	4	

- Molecule 53 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					AltConf
53	1	1	Total	C	N	O	P	0
			27	10	5	10	2	

- Molecule 54 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
54	1	2	Total	Mg	0
			2	2	

- Molecule 55 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).




Mol	Chain	Residues	Atoms					AltConf
55	1	1	Total	C	N	O	P	0
			31	10	5	13	3	


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 40S ribosomal protein SA




- Molecule 2: 40S ribosomal protein S3a




- Molecule 3: 40S ribosomal protein S2



- Molecule 4: 40S ribosomal protein S26




- Molecule 5: 40S ribosomal protein S4, X isoform







- Molecule 6: 60S ribosomal protein L41




- Molecule 7: 40S ribosomal protein S6




- Molecule 8: 40S ribosomal protein S7




- Molecule 9: 40S ribosomal protein S8



- Molecule 10: 40S ribosomal protein S9




- Molecule 11: 40S ribosomal protein S11




- Molecule 12: 40S ribosomal protein S13





- Molecule 13: 40S ribosomal protein S14




- Molecule 14: 40S ribosomal protein S21




- Molecule 15: 40S ribosomal protein S15a




- Molecule 16: 40S ribosomal protein S23




- Molecule 17: 40S ribosomal protein S24




- Molecule 18: 40S ribosomal protein S27




- Molecule 19: 40S ribosomal protein S30




- Molecule 20: 18S ribosomal RNA






• Molecule 21: 40S ribosomal protein S17




• Molecule 22: 40S ribosomal protein S16







- Molecule 23: 40S ribosomal protein S3




- Molecule 24: 40S ribosomal protein S5




- Molecule 25: 40S ribosomal protein S10




- Molecule 26: 40S ribosomal protein S12





- Molecule 27: 40S ribosomal protein S15







- Molecule 28: 40S ribosomal protein S18




- Molecule 29: 40S ribosomal protein S19




- Molecule 30: 40S ribosomal protein S20



- Molecule 31: 40S ribosomal protein S25




- Molecule 32: 40S ribosomal protein S28













- Molecule 33: 40S ribosomal protein S29









• Molecule 37: Eukaryotic translation initiation factor 3 subunit B





• Molecule 38: Eukaryotic translation initiation factor 3 subunit A









• Molecule 39: Eukaryotic translation initiation factor 3 subunit C








- Molecule 40: Eukaryotic translation initiation factor 3 subunit E




- Molecule 41: Eukaryotic translation initiation factor 3 subunit F






• Molecule 42: Eukaryotic translation initiation factor 3 subunit H




• Molecule 44: Eukaryotic translation initiation factor 3 subunit L


Chain L:




• Molecule 45: Eukaryotic translation initiation factor 3 subunit M

Chain M:









• Molecule 46: ATP-binding cassette sub-family E member 1




• Molecule 47: Eukaryotic translation initiation factor 1



• Molecule 48: Eukaryotic translation initiation factor 3 subunit D





ALA  
ASP  
TYR  
GLY  
GLY  
TYR  
ASP  
GLY  
GLY  
TYR  
VAL  
GLN  
ASP  
TYR  
GLU  
ASP  
PHE  
MET

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	5231	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	48	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.087	Depositor
Minimum map value	-0.033	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.004	Depositor
Recommended contour level	0.015	Depositor
Map size (Å)	423.6, 423.6, 423.6	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.059, 1.059, 1.059	Depositor



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, ZN, SF4, MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	a	0.59	0/1743	0.72	1/2367 (0.0%)
2	p	0.53	0/1743	0.71	1/2330 (0.0%)
3	d	0.62	0/1711	0.79	2/2310 (0.1%)
4	Q	0.66	0/829	0.74	0/1109
5	q	0.59	0/2074	0.74	1/2791 (0.0%)
6	W	0.49	0/232	0.58	0/294
7	r	0.48	0/1818	0.69	0/2421
8	s	0.54	0/1419	0.73	0/1895
9	t	0.55	0/1667	0.66	0/2223
10	c	0.57	0/1525	0.74	2/2035 (0.1%)
11	n	0.67	0/1140	0.66	0/1524
12	m	0.51	0/1227	0.67	0/1649
13	i	0.51	0/952	0.72	0/1275
14	y	0.57	0/632	0.65	0/844
15	f	0.63	1/1051 (0.1%)	0.76	1/1406 (0.1%)
16	j	0.58	0/1098	0.72	1/1464 (0.1%)
17	z	0.53	0/1017	0.67	0/1349
18	R	0.57	0/654	0.74	0/876
19	T	0.53	0/357	0.63	0/466
20	2	1.12	24/41041 (0.1%)	1.67	1229/63969 (1.9%)
21	w	0.50	0/1024	0.75	0/1377
22	g	0.44	0/1117	0.72	0/1494
23	b	0.44	0/1774	0.70	0/2387
24	e	0.43	0/1516	0.73	1/2037 (0.0%)
25	u	0.41	0/824	0.72	1/1111 (0.1%)
26	v	0.39	0/871	0.75	0/1168
27	o	0.43	0/999	0.73	1/1336 (0.1%)
28	k	0.39	0/1181	0.70	0/1581
29	x	0.40	0/1114	0.68	1/1493 (0.1%)
30	h	0.42	0/790	0.72	0/1059
31	P	0.38	0/564	0.74	1/758 (0.1%)
32	S	0.41	0/482	0.70	0/643

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
33	I	0.46	0/462	0.75	1/612 (0.2%)
34	U	0.43	0/475	0.70	0/626
35	V	0.43	0/2370	0.80	2/3221 (0.1%)
36	I	0.25	0/1495	0.52	0/2073
37	B	0.32	0/2981	0.64	0/4115
38	A	0.39	1/5468 (0.0%)	0.67	5/7401 (0.1%)
39	C	0.42	0/5157	0.72	3/6946 (0.0%)
40	E	0.39	0/3503	0.70	1/4728 (0.0%)
41	F	0.37	0/2126	0.72	3/2890 (0.1%)
42	H	0.41	0/2458	0.76	3/3313 (0.1%)
43	K	0.38	0/1785	0.75	4/2414 (0.2%)
44	L	0.41	0/3188	0.83	6/4299 (0.1%)
45	M	0.37	0/2757	0.71	2/3717 (0.1%)
46	I	0.44	1/4627 (0.0%)	0.74	4/6247 (0.1%)
47	N	0.40	0/739	0.77	1/989 (0.1%)
48	D	0.42	0/3700	0.74	5/5001 (0.1%)
49	Y	0.31	0/389	0.59	0/543
50	G	0.23	0/347	0.31	0/483
50	J	0.23	0/362	0.31	0/504
All	All	0.75	27/120575 (0.0%)	1.17	1283/171163 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	d	0	2
5	q	0	3
8	s	0	1
15	f	0	2
17	z	0	2
21	w	0	3
24	e	0	3
28	k	0	1
32	S	0	1
33	I	0	1
35	V	0	6
36	I	0	3
37	B	0	7
39	C	0	7
40	E	0	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	#Chirality outliers	#Planarity outliers
41	F	0	6
42	H	0	4
43	K	0	5
44	L	0	5
45	M	0	5
46	1	0	5
48	D	0	3
All	All	0	77

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
20	2	1100	A	N9-C4	-8.44	1.32	1.37
46	1	21	CYS	CB-SG	-6.79	1.70	1.82
15	f	93	LEU	C-N	-6.13	1.20	1.34
20	2	447	A	N9-C4	-6.13	1.34	1.37
20	2	640	A	N7-C5	-6.09	1.35	1.39

The worst 5 of 1283 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
20	2	1773	C	N1-C2-O2	16.15	128.59	118.90
20	2	1865	C	N3-C2-O2	-15.40	111.12	121.90
20	2	501	C	N1-C2-O2	15.07	127.94	118.90
20	2	1016	U	N3-C2-O2	-15.05	111.67	122.20
20	2	1865	C	C6-N1-C2	-14.94	114.32	120.30

There are no chirality outliers.

5 of 77 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	d	133	TYR	Peptide
3	d	135	GLY	Peptide
5	q	132	GLY	Peptide
5	q	184	THR	Peptide
5	q	87	MET	Peptide

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	a	214/216 (99%)	214 (100%)	0	0	100	100
2	p	209/211 (99%)	209 (100%)	0	0	100	100
3	d	214/216 (99%)	214 (100%)	0	0	100	100
4	Q	99/101 (98%)	99 (100%)	0	0	100	100
5	q	253/255 (99%)	250 (99%)	3 (1%)	0	100	100
6	W	22/24 (92%)	22 (100%)	0	0	100	100
7	r	220/222 (99%)	220 (100%)	0	0	100	100
8	s	165/181 (91%)	165 (100%)	0	0	100	100
9	t	195/205 (95%)	195 (100%)	0	0	100	100
10	c	178/180 (99%)	178 (100%)	0	0	100	100
11	n	131/144 (91%)	131 (100%)	0	0	100	100
12	m	147/149 (99%)	147 (100%)	0	0	100	100
13	i	123/125 (98%)	123 (100%)	0	0	100	100
14	y	80/82 (98%)	80 (100%)	0	0	100	100
15	f	127/129 (98%)	126 (99%)	1 (1%)	0	100	100
16	j	137/139 (99%)	135 (98%)	2 (2%)	0	100	100
17	z	120/122 (98%)	120 (100%)	0	0	100	100
18	R	80/82 (98%)	80 (100%)	0	0	100	100
19	T	40/44 (91%)	40 (100%)	0	0	100	100
21	w	124/130 (95%)	124 (100%)	0	0	100	100
22	g	136/138 (99%)	134 (98%)	2 (2%)	0	100	100
23	b	222/224 (99%)	221 (100%)	1 (0%)	0	100	100
24	e	187/189 (99%)	185 (99%)	2 (1%)	0	100	100
25	u	93/95 (98%)	93 (100%)	0	0	100	100
26	v	107/116 (92%)	107 (100%)	0	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
27	o	117/119 (98%)	117 (100%)	0	0	100	100
28	k	138/140 (99%)	138 (100%)	0	0	100	100
29	x	139/141 (99%)	139 (100%)	0	0	100	100
30	h	96/98 (98%)	96 (100%)	0	0	100	100
31	P	68/70 (97%)	68 (100%)	0	0	100	100
32	S	59/61 (97%)	59 (100%)	0	0	100	100
33	l	52/54 (96%)	52 (100%)	0	0	100	100
34	U	53/62 (86%)	53 (100%)	0	0	100	100
35	V	290/309 (94%)	287 (99%)	3 (1%)	0	100	100
36	I	301/316 (95%)	301 (100%)	0	0	100	100
37	B	528/566 (93%)	528 (100%)	0	0	100	100
38	A	682/722 (94%)	682 (100%)	0	0	100	100
39	C	615/743 (83%)	613 (100%)	2 (0%)	0	100	100
40	E	406/429 (95%)	406 (100%)	0	0	100	100
41	F	267/269 (99%)	266 (100%)	1 (0%)	0	100	100
42	H	289/318 (91%)	289 (100%)	0	0	100	100
43	K	215/217 (99%)	215 (100%)	0	0	100	100
44	L	370/372 (100%)	366 (99%)	4 (1%)	0	100	100
45	M	330/362 (91%)	330 (100%)	0	0	100	100
46	l	572/595 (96%)	570 (100%)	2 (0%)	0	100	100
47	N	89/91 (98%)	88 (99%)	1 (1%)	0	100	100
48	D	441/527 (84%)	441 (100%)	0	0	100	100
49	Y	76/78 (97%)	76 (100%)	0	0	100	100
50	G	68/258 (26%)	68 (100%)	0	0	100	100
50	J	71/258 (28%)	71 (100%)	0	0	100	100
All	All	9955/10894 (91%)	9931 (100%)	24 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM

entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	a	180/180 (100%)	175 (97%)	5 (3%)	43	68
2	p	192/192 (100%)	191 (100%)	1 (0%)	88	94
3	d	182/182 (100%)	179 (98%)	3 (2%)	62	79
4	Q	88/88 (100%)	88 (100%)	0	100	100
5	q	220/220 (100%)	219 (100%)	1 (0%)	88	94
6	W	23/23 (100%)	23 (100%)	0	100	100
7	r	193/193 (100%)	190 (98%)	3 (2%)	62	79
8	s	155/163 (95%)	154 (99%)	1 (1%)	86	92
9	t	174/178 (98%)	173 (99%)	1 (1%)	86	92
10	c	160/160 (100%)	160 (100%)	0	100	100
11	n	125/132 (95%)	124 (99%)	1 (1%)	81	89
12	m	130/130 (100%)	129 (99%)	1 (1%)	81	89
13	i	98/98 (100%)	98 (100%)	0	100	100
14	y	66/66 (100%)	64 (97%)	2 (3%)	41	66
15	f	112/112 (100%)	112 (100%)	0	100	100
16	j	111/111 (100%)	109 (98%)	2 (2%)	59	77
17	z	106/106 (100%)	106 (100%)	0	100	100
18	R	74/74 (100%)	73 (99%)	1 (1%)	67	81
19	T	35/35 (100%)	34 (97%)	1 (3%)	42	67
21	w	111/118 (94%)	109 (98%)	2 (2%)	59	77
22	g	114/114 (100%)	111 (97%)	3 (3%)	46	69
23	b	188/188 (100%)	186 (99%)	2 (1%)	73	85
24	e	159/159 (100%)	156 (98%)	3 (2%)	57	76
25	u	86/86 (100%)	85 (99%)	1 (1%)	71	84
26	v	94/98 (96%)	93 (99%)	1 (1%)	73	85
27	o	107/107 (100%)	106 (99%)	1 (1%)	78	88
28	k	122/122 (100%)	120 (98%)	2 (2%)	62	79
29	x	111/111 (100%)	109 (98%)	2 (2%)	59	77
30	h	91/91 (100%)	90 (99%)	1 (1%)	73	85

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
31	P	62/62 (100%)	62 (100%)	0	100	100
32	S	54/54 (100%)	54 (100%)	0	100	100
33	I	47/47 (100%)	47 (100%)	0	100	100
34	U	51/55 (93%)	50 (98%)	1 (2%)	55	75
35	V	256/268 (96%)	253 (99%)	3 (1%)	71	84
37	B	90/508 (18%)	90 (100%)	0	100	100
38	A	546/667 (82%)	541 (99%)	5 (1%)	78	88
39	C	554/662 (84%)	545 (98%)	9 (2%)	62	79
40	E	380/393 (97%)	376 (99%)	4 (1%)	73	85
41	F	237/237 (100%)	231 (98%)	6 (2%)	47	70
42	H	269/290 (93%)	265 (98%)	4 (2%)	65	81
43	K	192/192 (100%)	190 (99%)	2 (1%)	76	86
44	L	342/342 (100%)	337 (98%)	5 (2%)	65	81
45	M	305/324 (94%)	301 (99%)	4 (1%)	69	82
46	I	506/523 (97%)	498 (98%)	8 (2%)	62	79
47	N	77/78 (99%)	74 (96%)	3 (4%)	32	60
48	D	398/473 (84%)	389 (98%)	9 (2%)	50	72
All	All	7973/8812 (90%)	7869 (99%)	104 (1%)	70	82

5 of 104 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
39	C	726	LEU
42	H	133	LEU
48	D	368	ASP
40	E	136	LYS
41	F	230	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 110 such sidechains are listed below:

Mol	Chain	Res	Type
38	A	549	GLN
40	E	210	GLN
48	D	479	GLN
45	M	79	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
39	C	62	ASN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
20	2	1710/1720 (99%)	524 (30%)	10 (0%)

5 of 524 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
20	2	2	A
20	2	6	G
20	2	25	A
20	2	41	G
20	2	44	U

5 of 10 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
20	2	1474	A
20	2	1581	C
20	2	1807	C
20	2	550	C
20	2	912	C

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 5 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
52	SF4	1	601	-	0,12,12	-	-	-		
52	SF4	1	602	-	0,12,12	-	-	-		
53	ADP	1	603	54	24,29,29	0.95	1 (4%)	29,45,45	1.46	4 (13%)
55	ATP	1	605	54	26,33,33	0.93	1 (3%)	31,52,52	1.61	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
55	ATP	1	605	54	-	2/18/38/38	0/3/3/3
52	SF4	1	602	-	-	-	0/6/5/5
53	ADP	1	603	54	-	6/12/32/32	0/3/3/3
52	SF4	1	601	-	-	-	0/6/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
55	1	605	ATP	C5-C4	2.50	1.47	1.40
53	1	603	ADP	C5-C4	2.48	1.47	1.40

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
53	1	603	ADP	PA-O3A-PB	-3.72	120.07	132.83
55	1	605	ATP	PB-O3B-PG	-3.65	120.31	132.83
55	1	605	ATP	PA-O3A-PB	-3.49	120.84	132.83
55	1	605	ATP	C3'-C2'-C1'	3.42	106.13	100.98
55	1	605	ATP	N3-C2-N1	-3.16	123.74	128.68

There are no chirality outliers.


5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
53	1	603	ADP	C5'-O5'-PA-O1A
53	1	603	ADP	C5'-O5'-PA-O2A
55	1	605	ATP	O4'-C4'-C5'-O5'
53	1	603	ADP	PA-O3A-PB-O2B
53	1	603	ADP	PA-O3A-PB-O3B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
39	C	1
8	s	1
11	n	1
20	2	1
19	T	1
40	E	1
15	f	1

The worst 5 of 7 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	149:ASN	C	320:LYS	N	53.70
1	s	12:ASN	C	18:GLU	N	15.89
1	n	23:VAL	C	33:LEU	N	14.18
1	2	739:C	O3'	747:U	P	13.90
1	T	45:VAL	C	54:GLY	N	10.18


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-11458. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 166



Y Index: 218




Z Index: 202


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.015. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

## 6.6 Mask visualisation [i](#)


This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate [i](#)




The volume at the recommended contour level is 779 nm<sup>3</sup>; this corresponds to an approximate mass of 704 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.263 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.263 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.80	-	-
Author-provided FSC curve	3.76	6.74	3.93
Unmasked-calculated*	-	-	-

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-11458 and PDB model 6ZVJ. Per-residue inclusion information can be found in [section 3](#) on [page 16](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.015 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.015).







































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 66% of all backbone atoms, 63% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary ⓘ



































The table lists the average atom inclusion at the recommended contour level (0.015) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6280	 0.3650
1	 0.5520	 0.3700
2	 0.8920	 0.4510
A	 0.4590	 0.2820
B	 0.1580	 0.1870
C	 0.5260	 0.3210
D	 0.2240	 0.2230
E	 0.3830	 0.2210
F	 0.2050	 0.1790
G	 0.0230	 0.0940
H	 0.1980	 0.1700
I	 0.0010	 0.0250
J	 0.0050	 0.1020
K	 0.1220	 0.1540
L	 0.1220	 0.1220
M	 0.2460	 0.1820
N	 0.1230	 0.2020
P	 0.5120	 0.2920
Q	 0.7680	 0.4840
R	 0.7810	 0.4910
S	 0.5020	 0.3380
T	 0.7730	 0.4720
U	 0.4220	 0.2280
V	 0.5930	 0.3130
W	 0.6380	 0.4350
Y	 0.3330	 0.2290
a	 0.7940	 0.4990
b	 0.6250	 0.3870
c	 0.7990	 0.4990
d	 0.8000	 0.5040
e	 0.6180	 0.3680
f	 0.8380	 0.5370
g	 0.6820	 0.4140
h	 0.6050	 0.3840
i	 0.7780	 0.4680



*Continued on next page...*

*Continued from previous page...*

Chain	Atom inclusion	Q-score
j	 0.7930	 0.5160
k	 0.5290	 0.3190
l	 0.7530	 0.4120
m	 0.8010	 0.4880
n	 0.8240	 0.5230
o	 0.4410	 0.2630
p	 0.7850	 0.4830
q	 0.8290	 0.5100
r	 0.7600	 0.4440
s	 0.7230	 0.4310
t	 0.8090	 0.4720
u	 0.5380	 0.3070
v	 0.3850	 0.2290
w	 0.6980	 0.4320
x	 0.6420	 0.3460
y	 0.7940	 0.4940
z	 0.8030	 0.4900