



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 31, 2025 – 10:10 PM JST

PDB ID : 8ZX8 / pdb_00008zx8
Title : Structure-Based Mechanism and Specificity of Human Galactosyltransferase B3GalT5
Authors : Lo, J.M.; Ma, C.
Deposited on : 2024-06-14
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	278	85% 11% .
1	B	278	79% 16% .
2	E	2	100%
2	G	2	100%
3	F	2	100%
4	H	6	50% 50%
5	D	4	50% 50%
6	I	3	100%

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 4946 atoms, of which 119 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-1,3-galactosyltransferase 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	2	0
			2201	1427	371	390	13			
1	B	266	Total	C	N	O	S	0	0	0
			2189	1418	370	388	13			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	G	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	2	Total	C	N	O	0	0	0
			24	14	1	9			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	6	Total	C	H	N	O	0	0	0
			138	40	67	2	29			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	D	4	Total	C	N	O	0	0	0
			48	26	1	21			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-galactopyranose-(1-3)-alpha-D-galactopyranose-(1-4)-beta-D-galactopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	I	3	Total	C	N	O	0	0	0
			36	20	1	15			

- Molecule 7 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Mn	0	0
			1	1		
7	B	1	Total	Mn	0	0
			1	1		

- Molecule 8 is URIDINE-5'-DIPHOSPHATE (CCD ID: UDP) (formula: C₉H₁₄N₂O₁₂P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
8	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

- Molecule 9 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYLAMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: B3P) (formula: C₁₁H₂₆N₂O₆).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	A	1	Total	C	H	N	O	0	0
			45	11	26	2	6		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	B	1	Total	C	H	N	O	0	0
			45	11	26	2	6		

- Molecule 10 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			7	4	3		
10	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 11 is beta-D-galactopyranose (CCD ID: GAL) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 12 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	69	Total	O	0	0
			69	69		
12	B	18	Total	O	0	0
			18	18		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS failed to run properly.


- Molecule 1: Beta-1,3-galactosyltransferase 5

Chain A: 



- Molecule 1: Beta-1,3-galactosyltransferase 5

Chain B: 



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 



- Molecule 3: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1
FUC2

- Molecule 4: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-[α -L-fucopyranose-(1-6)]2-acetamido-2-deoxy- β -D-glucopyranose

Chain H:  50% 50%

MAG1
MAG2
BMA3
MAN4
MAN5
FUC6

- Molecule 5: 2-acetamido-2-deoxy- β -D-galactopyranose-(1-3)- α -D-galactopyranose-(1-4)- β -D-galactopyranose-(1-4)- β -D-glucopyranose

Chain D:  50% 50%

BGC1
GAL2
GLA3
NGA4

- Molecule 6: 2-acetamido-2-deoxy- β -D-galactopyranose-(1-3)- α -D-galactopyranose-(1-4)- β -D-galactopyranose

Chain I:  100%

GAL1
GLA2
NGA3

4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	47.80Å 86.53Å 87.64Å 90.00° 95.71° 90.00°	Depositor
Resolution (Å)	30.72 – 2.40	Depositor
% Data completeness (in resolution range)	85.8 (30.72-2.40)	Depositor
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, R_{free}	0.199 , 0.219	Depositor
Wilson B-factor (Å ²)	37.6	Xtriage
Anisotropy	0.162	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4946	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GLA, BMA, MN, GAL, UDP, B3P, PEG, FUC, NGA, MAN, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2266	0.73	0/3064
1	B	0.35	0/2248	0.58	0/3040
All	All	0.43	0/4514	0.66	0/6104

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	209	ARG	Sidechain
1	A	84	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2201	0	2194	18	0
1	B	2189	0	2175	33	0
2	E	28	0	25	0	0
2	G	28	0	25	1	0
3	F	24	0	22	0	0
4	H	71	67	61	0	0
5	D	48	0	42	4	0
6	I	36	0	31	1	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
8	A	25	0	11	0	0
8	B	25	0	11	0	0
9	A	19	26	26	0	0
9	B	19	26	26	0	0
10	A	14	0	20	1	0
11	A	11	0	10	2	0
12	A	69	0	0	0	0
12	B	18	0	0	0	0
All	All	4827	119	4679	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:A:406:GAL:C1	5:D:4:NGA:O3	2.19	0.91
5:D:3:GLA:C3	5:D:4:NGA:C1	2.47	0.90
1:A:232:VAL:HG21	1:A:252:ARG:HG2	1.64	0.80
5:D:3:GLA:O3	5:D:4:NGA:C2	2.30	0.80
1:B:154:LYS:O	1:B:218:GLY:HA2	1.86	0.75
1:A:176:THR:HG23	1:A:177:THR:HG23	1.70	0.74
1:B:45:LEU:HD11	1:B:123:ASP:HB2	1.75	0.69
1:A:232:VAL:CG2	1:A:252:ARG:HG2	2.23	0.68
1:A:80:TRP:CZ3	1:A:161[B]:ILE:HD12	2.37	0.60
1:B:45:LEU:HG	1:B:122:LYS:HA	1.84	0.59
1:B:288:LYS:HE3	1:B:290:ARG:HB3	1.84	0.59
1:B:232:VAL:HG21	1:B:252:ARG:HG2	1.85	0.58
1:B:124:PHE:HA	2:G:1:NAG:H62	1.85	0.58
1:B:241:LEU:HD13	6:I:3:NGA:H62	1.87	0.56
1:A:172:LYS:O	1:A:172:LYS:HG2	2.05	0.56
1:A:190:PRO:HD2	1:A:208:ASP:O	2.07	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:PRO:HB3	1:A:148:GLN:HB3	1.91	0.52
1:B:165:TYR:CE2	1:B:169:LEU:HD11	2.45	0.52
1:B:66:SER:O	1:B:73:ARG:NH1	2.38	0.51
1:B:153:MET:HB2	1:B:220:VAL:HG22	1.92	0.51
5:D:3:GLA:H3	5:D:4:NGA:C1	2.40	0.51
1:B:45:LEU:HD11	1:B:123:ASP:CB	2.41	0.50
1:B:232:VAL:CG2	1:B:252:ARG:HG2	2.41	0.50
1:B:154:LYS:O	1:B:218:GLY:CA	2.58	0.50
1:B:217:THR:HG21	1:B:268:PRO:HB3	1.92	0.50
1:A:58:PHE:CB	1:A:89:LYS:HB3	2.42	0.50
1:B:197:LYS:HG3	1:B:198:TRP:CD1	2.47	0.49
1:B:193:GLN:HB3	1:B:195:PHE:CE2	2.47	0.48
1:B:73:ARG:HG2	1:B:95:PHE:CD1	2.48	0.48
1:B:58:PHE:CB	1:B:89:LYS:HB3	2.42	0.48
1:B:154:LYS:HD2	1:B:219:TYR:CZ	2.48	0.48
1:A:61:LEU:HD23	1:A:153:MET:HB3	1.94	0.48
1:B:187:ASN:OD1	1:B:209:ARG:NH2	2.42	0.48
1:B:278:PHE:HA	1:B:281:ILE:HG12	1.96	0.48
1:B:45:LEU:HD11	1:B:123:ASP:CA	2.45	0.47
1:A:232:VAL:HG21	1:A:252:ARG:CG	2.41	0.46
1:B:67:HIS:ND1	1:B:109:VAL:HG21	2.30	0.46
1:A:76:ILE:HG12	1:A:80:TRP:CZ2	2.51	0.45
1:A:307:CYS:O	1:A:308:PRO:C	2.53	0.45
1:B:273:PHE:HB2	1:B:295:TYR:CZ	2.52	0.45
1:B:232:VAL:HG21	1:B:252:ARG:CG	2.47	0.44
1:B:175:ARG:CZ	1:B:175:ARG:HB3	2.47	0.44
1:B:216:GLY:O	1:B:217:THR:C	2.56	0.44
1:B:43:ASN:HB3	1:B:44:PHE:H	1.49	0.43
1:B:285:HIS:CD2	1:B:286:PHE:H	2.36	0.43
1:A:211:PRO:HG3	1:A:248:LEU:HD23	2.01	0.43
1:B:58:PHE:HB3	1:B:89:LYS:HB3	2.00	0.43
1:A:47:LEU:H	10:A:404:PEG:H42	1.83	0.42
1:B:131:LEU:HD23	1:B:131:LEU:HA	1.93	0.42
1:B:44:PHE:HA	1:B:122:LYS:HG2	2.00	0.42
11:A:406:GAL:C1	11:A:406:GAL:O4	2.68	0.41
1:A:63:VAL:HG22	1:A:155:THR:CG2	2.50	0.41
1:A:185:LYS:HB3	1:A:188:GLU:HG3	2.02	0.41
1:B:180:PHE:CD2	1:B:221:PHE:HB3	2.56	0.41
1:A:128:TYR:CZ	1:A:197:LYS:HD3	2.57	0.40
1:A:154:LYS:HD2	1:A:219:TYR:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/278 (96%)	261 (98%)	6 (2%)	0	100	100
1	B	264/278 (95%)	257 (97%)	7 (3%)	0	100	100
All	All	531/556 (96%)	518 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/255 (96%)	246 (100%)	0	100	100
1	B	244/255 (96%)	243 (100%)	1 (0%)	89	95
All	All	490/510 (96%)	489 (100%)	1 (0%)	92	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	172	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	231	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

19 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	BGC	D	1	5	12,12,12	0.15	0	17,17,17	0.42	0
5	GAL	D	2	5	11,11,12	0.38	0	15,15,17	0.66	0
5	GLA	D	3	5	11,11,12	0.28	0	15,15,17	0.61	0
5	NGA	D	4	5	14,14,15	0.43	0	17,19,21	0.74	0
2	NAG	E	1	1,2	14,14,15	0.29	0	17,19,21	0.49	0
2	NAG	E	2	2	14,14,15	0.28	0	17,19,21	0.56	0
3	NAG	F	1	1,3	14,14,15	0.42	0	17,19,21	0.73	0
3	FUC	F	2	3	10,10,11	0.32	0	14,14,16	0.49	0
2	NAG	G	1	1,2	14,14,15	0.44	0	17,19,21	0.55	0
2	NAG	G	2	2	14,14,15	0.41	0	17,19,21	0.76	1 (5%)
4	NAG	H	1	4,1	14,14,15	0.44	0	17,19,21	1.29	2 (11%)
4	NAG	H	2	4	14,14,15	0.45	0	17,19,21	0.75	1 (5%)
4	BMA	H	3	4	11,11,12	0.68	0	15,15,17	0.63	0
4	MAN	H	4	4	11,11,12	0.46	0	15,15,17	0.62	0
4	MAN	H	5	4	11,11,12	0.31	0	15,15,17	0.80	0
4	FUC	H	6	4	10,10,11	0.34	0	14,14,16	0.61	1 (7%)
6	GAL	I	1	6	11,11,12	0.77	1 (9%)	15,15,17	2.17	4 (26%)
6	GLA	I	2	6	11,11,12	1.74	2 (18%)	15,15,17	0.70	0
6	NGA	I	3	6	14,14,15	0.40	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BGC	D	1	5	-	0/2/22/22	0/1/1/1
5	GAL	D	2	5	-	2/2/19/22	0/1/1/1
5	GLA	D	3	5	-	2/2/19/22	0/1/1/1
5	NGA	D	4	5	-	0/6/23/26	0/1/1/1
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
3	NAG	F	1	1,3	-	2/6/23/26	0/1/1/1
3	FUC	F	2	3	-	-	0/1/1/1
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	5/6/23/26	0/1/1/1
4	NAG	H	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	H	2	4	-	0/6/23/26	0/1/1/1
4	BMA	H	3	4	-	0/2/19/22	0/1/1/1
4	MAN	H	4	4	-	0/2/19/22	0/1/1/1
4	MAN	H	5	4	-	1/2/19/22	0/1/1/1
4	FUC	H	6	4	-	-	0/1/1/1
6	GAL	I	1	6	-	2/2/19/22	0/1/1/1
6	GLA	I	2	6	-	0/2/19/22	0/1/1/1
6	NGA	I	3	6	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	I	2	GLA	O5-C1	4.66	1.51	1.43
6	I	2	GLA	O5-C5	2.38	1.48	1.43
6	I	1	GAL	O5-C5	2.10	1.47	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	I	1	GAL	C1-O5-C5	6.53	121.04	112.19
6	I	1	GAL	O5-C5-C6	-3.13	102.29	107.20
4	H	1	NAG	C2-N2-C7	2.95	127.11	122.90
6	I	1	GAL	C2-C3-C4	-2.54	106.50	110.89
4	H	1	NAG	C1-O5-C5	2.50	115.58	112.19
2	G	2	NAG	C1-O5-C5	2.47	115.53	112.19
6	I	1	GAL	O5-C5-C4	2.34	116.53	110.83
4	H	2	NAG	C1-C2-N2	2.10	114.08	110.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	6	FUC	C1-C2-C3	2.09	112.24	109.67

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	G	2	NAG	C3-C2-N2-C7
2	G	2	NAG	C8-C7-N2-C2
2	G	2	NAG	O7-C7-N2-C2
5	D	3	GLA	O5-C5-C6-O6
3	F	1	NAG	C8-C7-N2-C2
4	H	1	NAG	C8-C7-N2-C2
5	D	2	GAL	O5-C5-C6-O6
5	D	3	GLA	C4-C5-C6-O6
5	D	2	GAL	C4-C5-C6-O6
6	I	1	GAL	O5-C5-C6-O6
3	F	1	NAG	O7-C7-N2-C2
4	H	1	NAG	O7-C7-N2-C2
6	I	1	GAL	C4-C5-C6-O6
2	G	2	NAG	O5-C5-C6-O6
2	G	2	NAG	C1-C2-N2-C7
4	H	5	MAN	O5-C5-C6-O6


There are no ring outliers.

4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	4	NGA	4	0
6	I	3	NGA	1	0
2	G	1	NAG	1	0
5	D	3	GLA	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







Oligosaccharide Chain H



Oligosaccharide Chain D





5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
11	GAL	A	406	-	11,11,12	0.73	0	15,15,17	2.04	2 (13%)
10	PEG	A	405	-	6,6,6	0.11	0	5,5,5	0.10	0
9	B3P	A	403	-	18,18,18	0.56	0	21,23,23	1.02	1 (4%)
8	UDP	A	402	7	24,26,26	0.53	0	37,40,40	0.50	0
9	B3P	B	403	-	18,18,18	0.68	0	21,23,23	1.52	4 (19%)
10	PEG	A	404	-	6,6,6	0.10	0	5,5,5	0.09	0
8	UDP	B	402	7	24,26,26	0.52	0	37,40,40	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	GAL	A	406	-	-	2/2/19/22	0/1/1/1
10	PEG	A	405	-	-	3/4/4/4	-
9	B3P	A	403	-	-	0/28/28/28	-
8	UDP	A	402	7	-	3/16/32/32	0/2/2/2
9	B3P	B	403	-	-	0/28/28/28	-
10	PEG	A	404	-	-	2/4/4/4	-
8	UDP	B	402	7	-	7/16/32/32	0/2/2/2

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	A	406	GAL	C1-O5-C5	6.21	120.61	112.19
11	A	406	GAL	O5-C1-C2	4.08	117.08	110.77
9	B	403	B3P	O6-C7-C4	-3.48	104.58	111.63
9	B	403	B3P	C2-N2-C8	-2.84	112.05	116.08
9	B	403	B3P	O1-C9-C8	-2.83	105.91	111.63
9	B	403	B3P	C11-C8-C9	2.37	115.06	110.04
9	A	403	B3P	C7-C4-C5	-2.26	105.26	110.04

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	402	UDP	PA-O3A-PB-O3B
8	B	402	UDP	C5'-O5'-PA-O1A
8	B	402	UDP	O4'-C4'-C5'-O5'
11	A	406	GAL	O5-C5-C6-O6
10	A	405	PEG	O2-C3-C4-O4
10	A	404	PEG	C1-C2-O2-C3
10	A	405	PEG	O1-C1-C2-O2
10	A	404	PEG	O2-C3-C4-O4
11	A	406	GAL	C4-C5-C6-O6
8	B	402	UDP	PB-O3A-PA-O5'
8	B	402	UDP	C5'-O5'-PA-O3A
10	A	405	PEG	C1-C2-O2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
8	B	402	UDP	C5'-O5'-PA-O2A
8	B	402	UDP	C3'-C4'-C5'-O5'
8	A	402	UDP	PB-O3A-PA-O1A
8	B	402	UDP	C4'-C5'-O5'-PA
8	A	402	UDP	PA-O3A-PB-O1B


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	A	406	GAL	2	0
10	A	404	PEG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand GAL A 406



Ligand B3P A 403







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.


6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

6.3 Carbohydrates [i](#)




EDS failed to run properly - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








6.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UDP A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UDP B 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around B3P A 403:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around B3P B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.